



Index to Volume 121

The Auk 121(4):1306–1334, 2004

COMPILED BY TODD J. UNDERWOOD, AMBER J. ROBINSON, AND C. JOAN PATTERSON

The index is confined primarily to new information. New specific names are printed in **boldface** type.

100 Years Ago in The American Ornithologists'

Union, 278–279, 629–630, 984, 1302–1303
2% rule, 1–6
abundance, 865–876
Aburria aburri, 682–694
Acanthis flammea, 1238–1249
Acanthisitta, 435–445
A. chloris, 1155–1160
Acanthisittidae, 1155–1160
Acanthiza pusilla, 1199–1207
Acanthochaera inauris, 1155–1160
Accentor, Alpine, 184–191
Accipiter bicolor, 515–525
A. cooperii, 738–751, 904–916
A. melanoleucus, 1226–1237
A. nisus, 1103–1121
A. striatus, 738–751, 837–846, 904–916
A. superciliosus, 88–102
A. tachiro, 1226–1237
Accipitridae, 88–102
Acrocephalus arundinaceus, 203–209, 435–445, 1226–1237
A. palustris, 203–209
A. schoenobaenus, 203–209, 1155–1160
A. scirpaceus, 203–209
Actitis hypoleucos, 985–995
A. macularia, 391–403, 985–995
A. macularius, 985–995
activity, — budgets, 1161–1171; — patterns, 380–390
Actophilornis, 391–403
A. africanus, 391–403
Adair, Bill, see Dove, Carla J., et al.
Adamo, Claudia, see Puglisi, Luca, and —
Adams, Lowell W., In Memoriam: Daniel Loney Leedy, 1912–2003, 972
adaptive developmental plasticity, 1187–1198
adrenocortical response, 1094–1102, 1238–1249
Aegithalidae, 1155–1160
Aegithalos caudatus, 777–787, 1155–1160
A. concinnus, 1155–1160
Aegithina tithia, 1155–1160
Aegolius funereus, 1103–1121
Aepyodius arfakianus, 308–317
aerodynamics, 973–976
Aethia cristatella, 452–462

Afton, Alan D., see Anteau, Michael J., and —
Agelaius humeralis, 1155–1160
A. phoeniceus, 184–191, 224–229, 361–371, 435–445, 446–451, 767–776, 824–836, 877–893, 1081–1093, 1146–1154, 1226–1237
aging nests, 134–147
agricultural, — land, 361–371; — landscape, 404–414, 726–737, 1146–1154
Aguarua salicifolia, 23–34
Aimophila nivalis, 74–87
A. cassinii, 1081–1093
A. ruficeps canescens, 904–916
A. ruficeps, 446–451, 904–916
Aix sponsa, 148–155
Akalat, Rubeho, 23–34
Aki trees, 548–556
Akiapolaau, 548–556
Akikiki, 1–6
Alavahio, Maui, 1–6
Alauda arvensis, 796–805, 1155–1160
Alaudidae, 1155–1160
Albatross, 422–434, 847–864; Black-browed, 463–472; Gray-headed, 1208–1213; Wandering, 1208–1213; Yellow-nosed, 463–472
Albrecht, Tomas, Edge effect in wetland-arable land boundary determines nesting success of Scarlet Rosefinches (*Carpodacus erythrinus*) in the Czech Republic, 361–371
Alca, 452–462
A. torda, 452–462
Alcedo guttata, 985–995
Alcidae, 35–45, 148–155, 452–462
alcids, 35–45, 148–155, 452–462, 1161–1171
Alectoris, 1–6
Alectura lathamii, 308–317
Alethe, 23–34
Alisauskas, Ray T., see Traylor, Joshua J., et al.
alpha-2-macroglobulin levels, 717–725
altitudinal movements, 1269–1281
Altshuler, Douglas L., see Stiles, F. Gary, and —
Amakihi, 1–6; Hawaii, 548–556
Amandava subflava, 156–161
Amazilia, 985–995
A. amabilis, 985–995
A. decora, 985–995



Index to Volume 121

The Auk 121(4):1306–1334, 2004

COMPILED BY TODD J. UNDERWOOD, AMBER J. ROBINSON, AND C. JOAN PATTERSON

The index is confined primarily to new information. New specific names are printed in **boldface** type.

100 Years Ago in The American Ornithologists'

Union, 278–279, 629–630, 984, 1302–1303
2% rule, 1–6
abundance, 865–876
Aburria aburri, 682–694
Acanthis flammea, 1238–1249
Acanthisitta, 435–445
A. chloris, 1155–1160
Acanthisittidae, 1155–1160
Acanthiza pusilla, 1199–1207
Acanthochaera inauris, 1155–1160
Accentor, Alpine, 184–191
Accipiter bicolor, 515–525
A. cooperii, 738–751, 904–916
A. melanoleucus, 1226–1237
A. nisus, 1103–1121
A. striatus, 738–751, 837–846, 904–916
A. superciliosus, 88–102
A. tachiro, 1226–1237
Accipitridae, 88–102
Acrocephalus arundinaceus, 203–209, 435–445, 1226–1237
A. palustris, 203–209
A. schoenobaenus, 203–209, 1155–1160
A. scirpaceus, 203–209
Actitis hypoleucos, 985–995
A. macularia, 391–403, 985–995
A. macularius, 985–995
activity, — budgets, 1161–1171; — patterns, 380–390
Actophilornis, 391–403
A. africanus, 391–403
Adair, Bill, see Dove, Carla J., et al.
Adamo, Claudia, see Puglisi, Luca, and —
Adams, Lowell W., In Memoriam: Daniel Loney Leedy, 1912–2003, 972
adaptive developmental plasticity, 1187–1198
adrenocortical response, 1094–1102, 1238–1249
Aegithalidae, 1155–1160
Aegithalos caudatus, 777–787, 1155–1160
A. concinnus, 1155–1160
Aegithina tithia, 1155–1160
Aegolius funereus, 1103–1121
Aepyodius arfakianus, 308–317
aerodynamics, 973–976
Aethia cristatella, 452–462

Afton, Alan D., see Anteau, Michael J., and —
Agelaius humeralis, 1155–1160
A. phoeniceus, 184–191, 224–229, 361–371, 435–445, 446–451, 767–776, 824–836, 877–893, 1081–1093, 1146–1154, 1226–1237
aging nests, 134–147
agricultural, — land, 361–371; — landscape, 404–414, 726–737, 1146–1154
Aguarua salicifolia, 23–34
Aimophila nestivalis, 74–87
A. cassinii, 1081–1093
A. ruficeps canescens, 904–916
A. ruficeps, 446–451, 904–916
Aix sponsa, 148–155
Akalat, Rubeho, 23–34
Aki trees, 548–556
Akiapolaau, 548–556
Akikiki, 1–6
Alavahio, Maui, 1–6
Alauda arvensis, 796–805, 1155–1160
Alaudidae, 1155–1160
Albatross, 422–434, 847–864; Black-browed, 463–472; Gray-headed, 1208–1213; Wandering, 1208–1213; Yellow-nosed, 463–472
Albrecht, Tomas, Edge effect in wetland-arable land boundary determines nesting success of Scarlet Rosefinches (*Carpodacus erythrinus*) in the Czech Republic, 361–371
Alca, 452–462
A. torda, 452–462
Alcedo guttata, 985–995
Alcidae, 35–45, 148–155, 452–462
alcids, 35–45, 148–155, 452–462, 1161–1171
Alectoris, 1–6
Alectura lathamii, 308–317
Alethe, 23–34
Alisauskas, Ray T., see Traylor, Joshua J., et al.
alpha-2-macroglobulin levels, 717–725
altitudinal movements, 1269–1281
Altshuler, Douglas L., see Stiles, F. Gary, and —
Amakihi, 1–6; Hawaii, 548–556
Amandava subflava, 156–161
Amazilia, 985–995
A. amabilis, 985–995
A. decora, 985–995

- Amazon, Yellow-crowned, 318-332; Yellow-headed, 318-332; Yellow-naped, 318-332
Amazona, 230-242, 318-332, 1214-1225
A. aestiva, 230-242, 318-332
A. amazonica, 318-332
A. auropalliata, 318-332
A. a. auropalliata, 318-332
A. a. caribaea, 318-332
A. a. parvipes, 318-332
A. autumnalis, 88-102, 318-332
A. barbadensis, 318-332
A. farinosa, 88-102, 318-332
A. finschi, 1214-1225
A. ochrocephala, 318-332
A. o. auropalliata, 230-242
A. o. nattereri, 318-332
A. o. ochrocephala, 318-332
A. o. panamensis, 318-332
A. o. xantholaema, 318-332
A. oratrix, 318-332
A. o. belizensis, 318-332
A. o. guatemalensis, 318-332
A. o. hondurensis, 318-332
A. o. magna, 318-332
A. o. oratrix, 318-332
A. o. tresmariae, 318-332
A. viridigenalis, 318-332, 1214-1225
Amblycercus holosericeus, 224-229
Ammodramus, 1081-1093
A. bairdii, 1130-1145
A. maritimus, 985-995
A. savannarum, 435-445, 1130-1145
Amphispiza bilineata, 1081-1093
 Amrhein, Valentin, et al., Seasonal patterns of singing activity vary with time of day in the Nightingale (*Luscinia megarhynchos*), 110-117
Anabacerthia, 752-766
Anas, 695-706
A. acuta, 497-508, 587-599, 695-706
A. americana, 587-599
A. clypeata, 497-508, 587-599, 695-706
A. crecca, 587-599
A. diazi, 930-949
A. discors, 497-508, 526-540, 587-599, 695-706
A. fulvigula, 930-949
A. luzonica, 930-949
A. platyrhynchos, 497-508, 526-540, 587-599, 695-706, 917-929, 930-949
A. poecilorhyncha, 930-949
A. p. haringtoni, 930-949
A. p. poecilorhyncha, 930-949
A. p. zonorhyncha, 930-949
A. rubripes, 930-949
A. strepera, 497-508, 587-599, 695-706, 950-962
A. superciliosa, 930-949
A. wyvilliana, 930-949
A. zonorhyncha, 930-949
Anatidae, 930-949
 ancestral polymorphism, 930-949
 Anderson, Mark D., see Covas, Rita, et al.
 Andersson, Staffan, review by, 267-269
Andropadus, 23-34
A. latirostris, 435-445
 aningas, 35-45
 Anhingidae, 35-45
Anodorhynchus, 230-242
A. hyacinthinus, 230-242
A. leari, 230-242
Anser, 1-6
A. caerulescens caerulescens, 652-655
A. hutchinsii, 985-995
 Anseriformes, 35-45, 354-360, 682-694
 ant chemicals, 1262-1268
 Antshrike, Acre, 1031-1039; Amazonian, 1031-1039; Barred, 1031-1039; Streak-backed, 1031-1039
 Antarctic storm, 1208-1213
 Antbird, 88-102; Dusky, 486-496; White-bellied, 486-496
 Anteau, Michael J., and Afton, Alan D., Nutrient reserves of Lesser Scaup (*Aythya affinis*) during spring migration in the Mississippi Flyway: A test of the spring condition hypothesis, 917-929
Anthracothorax nigricollis, 88-102
Anthreptes singalensis, 1155-1160
Anthus pratensis, 1155-1160
A. richardi, 35-45
A. rubescens, 1081-1093, 1250-1261
A. spinoletta, 777-787
A. spragueii, 1081-1093, 1130-1145
 anting behaviour, 1262-1268
 Antwren, Checker-throated, 486-496; Moustached, 985-995; Pygmy, 985-995; Pygmy complex, 985-995
Aphelocoma californica, 1081-1093
Aphriza virgata, 752-766
Aplonis panayensis, 435-445
Apodidae, 88-102, 1187-1198
Aptenodytes, 422-434
A. forsteri, 422-434
A. patagonicus, 422-434, 1161-1171
Apteryx, 308-317
Apus apus, 1187-1198
A. cafer, 985-995
A. pacificus, 985-995
Aquila rapax, 1081-1093
Ara, 230-242
A. ambigua, 985-995
A. ambiguus, 985-995
A. ararauna, 88-102, 230-242
A. chloroptera, 88-102
A. erythrocephala, 985-995
A. erythrocephalus, 985-995
A. erythrura, 985-995
A. erythrurus, 985-995
A. macao, 88-102
A. manilata, 88-102

- A. militaris*, 985–995
A. severa, 230–242, 985–995
A. severus, 985–995
Arachnothera, 435–445
A. flavigaster, 435–445
Aratinga, 230–242
A. aurea, 230–242
A. cactorum, 230–242
A. finschi, 985–995
A. leucophthalma, 985–995
Arenaria interpres interpres, 333–344
Archilochus colubris, 877–893, 1040–1055
 Arctic breeders, 1238–1249
Ardeola ralloides, 192–202
 area, — requirements, 58–67; — sensitivity, 767–776, 1130–1145
 area-sensitive distribution pattern, 904–916
Arenaria interpres interpres, 333–344
 Arini, 230–242
 Armesto, Juan J., see Reid, Sharon, et al.
 arrival date, 1269–1281
Artamus pelewensis, 1155–1160
 arthropod biomass, 877–893
 artificial nests, 777–787
 Asio, 35–45
 asymmetry, 473–485
Atrichornis, 1155–1160
Attila spadiceus, 88–102
 Auk, Great, 35–45
 Auklet, Cassin's, 452–462; Crested, 452–462;
 Rhinceros, 452–462, 463–472
Aulacorhynchus prasinus, 985–995
 Austin, Jeremy J., et al., A global molecular
 phylogeny of the small *Puffinus* shearwaters
 and implications for systematics of the Little-
 Audubon's Shearwater complex, 847–864
 austral system, 1005–1013
 Avise, John C., Molecular Markers, Natural History,
 and Evolution, 2nd ed. (rev.), 1298–1299
Aythya affinis, 587–599, 917–929
A. marila, 917–929
A. valisineria, 170–177, 333–344

Bacillus licheniformis, 1262–1268; feather-degrading,
 652–655, 656–659
 bacterial degradation, 652–655, 656–659
 bactericidal activity, 1262–1268
Baeolophus bicolor, 74–87, 435–445, 738–751, 865–876
 Bairlein, Franz, and Winkel, Wolfgang, In Memoriam:
 Friedrich Walter Goethe, 1911–2003, 1292
 Baker, Allan J., see Pereira, Sérgio Luiz, and —
 Ball, Tim, see Houston, Stuart, et al.
 bamboo, 515–525
 Bananaquit, 652–655
 Banks, Richard C., et al., Forty-fifth Supplement to
 the American Ornithologists' Union Check-list of
 North American Birds, 985–995
 barbets, 88–102, 178–183
 Barg, Jennifer J., see Jones, Jason, et al.
Basilinna, 985–995
 Batts, H. Lewis, Jr., 1922–2001, In Memoriam, 970
 Bee-eater, 811–823, 1122–1129; Blue-throated,
 811–823; Carmine, 811–823; European, 541–547,
 811–823; Little, 811–823; Little Green, 811–823;
 Rainbow, 811–823; Red-throated, 811–823;
 White-fronted, 811–823
 behavior, 565–586; clustering, 877–893; feeding,
 509–514; foraging, 210–223; foray, 486–496;
 incidental, 1161–1171; loafing, 380–390;
 reproductive, 1070–1080; spacing, 58–67;
 territorial, 486–496
 Beissinger, Steven R., see Sheridan, Jennifer A., et al.
 Benhamou, Simon, see Tourenq, Christophe, et al.
 Beresford, Pamela, et al., A new species of akalat
 (*Sheppardia*) narrowly endemic in the Eastern
 Arc of Tanzania, 23–34
 Bermingham, Eldredge, see Eberhard, Jessica R.,
 and —
 Bertelli, Sara, see Giannini, Norberto P., and —
 bill length, 178–183
 biogeography, 318–332, 637–651, 660–681, 752–766,
 837–846
 biological diversification, 1014–1018
 Bird, Tussock, 752–766
 birds of paradise, 637–651
 birds of prey, 894–903
 Bishop, Red, 1226–1237
 Bittern, Great, 541–547
 Blackbird, 224–229; Red-winged, 184–191, 361–371,
 446–451, 767–776, 824–836, 877–893, 1081–1093,
 1146–1154, 1226–1237; Rusty, 162–169; Yellow-
 headed, 1226–1237
 Black-Tyrant, White-winged, 1005–1013
Bleda, 23–34
 blood, — parasites, 837–846; — sampling, 354–360
 Bluebird, Eastern, 118–133, 302–307, 652–655;
 Mountain, 118–133, 1081–1093; Western, 118–133
 Bobolink, 767–776, 1103–1121, 1250–1261
 Bock, Walter J., Ernst Mayr at 100: A life inside and
 outside of ornithology, 637–651
 body, — condition, 170–177, 917–929; — size, 68–73;
 — stores, 333–344
 Boersma, P. Dee, et al., Why penguin eggshells are
 thick, 148–155
 Boland, Christopher R. J., Breeding biology of
 Rainbow Bee-eaters (*Merops ornatus*): A
 migratory, colonial, cooperative bird, 811–823
 Bolger, Douglas T., see Morrison, Scott A., et al.
 Bollinger, Eric K., and Gavin, Thomas A., Responses
 of nesting Bobolinks (*Dolichonyx oryzivorus*) to
 habitat edges, 767–776
 Bollinger, Eric K., see Walk, Jeffery W., et al.
Bombicilla cedrorum, 435–445
B. garrulus, 1155–1160
 Bombycillidae, 1155–1160
 booms, 541–547

- Borgmann, Kathi L., et al., Wintering Yellow-rumped Warblers (*Dendroica coronata*) track manipulated abundance of *Myrica cerifera* fruits, 74–87
- Botaurus stellaris*, 541–547
- Bowie, Rauri C. K., et al., Systematics and biogeography of double-collared sunbirds from the Eastern Arc Mountains, Tanzania, 660–681
- Bowlin, Melissa S., and Winkler, David W., Natural variation in flight performance is related to timing of breeding in Tree Swallows (*Tachycineta bicolor*) in New York, 345–353
- Bowman, Jacob L., see Perkins, Kelly A., et al.
- Brachyramphus*, 985–995
- B. brevirostris*, 35–45
- B. marmoratus*, 1161–1171
- Brambling, 1238–1249
- Brant, Black, 68–73, 210–223
- Branta*, 1–6
- B. bernicla*, 210–223, 985–995
- B. b. nigricans*, 68–73, 210–223
- B. canadensis*, 985–995
- B. c. asiatica*, 985–995
- B. c. canadensis*, 985–995
- B. c. fulva*, 985–995
- B. c. hutchinsii*, 985–995
- B. c. interior*, 985–995
- B. c. leucopareia*, 985–995
- B. c. maxima*, 985–995
- B. c. minima*, 985–995
- B. c. moffitti*, 985–995
- B. c. occidentalis*, 985–995
- B. c. parvipes*, 985–995
- B. c. taverneri*, 985–995
- B. hutchinsii*, 985–995
- B. leucopsis*, 68–73, 985–995
- B. sandvicensis*, 985–995
- Breckenridge, Walter John, 1903–2003, In Memoriam, 1286–1288
- breeders, capital, 333–344; income, 333–344
- breeding, — biology, 565–586, 811–823, 1226–1237; — characters, 422–434; — cycle, 110–117; — density, 302–307; — dispersion, 811–823; — grounds, high-Arctic, 333–344; — habitat, 726–737; — performance, 1103–1121; — population, 68–73; — season, 58–67; — sites, 257–262; — synchrony, 302–307
- breeding-season decisions, 796–805
- Bretagnolle, Vincent, see Austin, Jeremy J., et al.
- Breuner, Creagh W., see Hahn, Thomas P., et al.
- Brewster Memorial Award, 2003: Douglas W. Mock, 264–265
- Brittingham, Margaret C., see Rodewald, Paul G., and —
- broadbills, 1155–1160
- brood parasitism, 156–161, 1122–1129, 1172–1186; conspecific, 1122–1129
- brood, — reduction, 1214–1225; — size, 1070–1080
- Brotogeris chrysopterus*, 88–102
- Brown, Charles R., see Covas, Rita, et al.
- Brown, Charles R., see Weaver, Heath B., and —
- Brown, Mary Bomberger, see Covas, Rita, et al.
- Browne, Beth Ann, see Goldstein, Gerald, et al.
- Brumfield, Robb T., see Whitney, Bret M., et al.
- Brush-Finch, Stripe-headed, 289–301
- Brush-turkey, Australian, 308–317; Wattled, 308–317
- Brzęk, Paweł, and Konarzewski, Marek, Effect of refeeding on growth, development, and behavior of undernourished Bank Swallow (*Riparia riparia*) nestlings, 1187–1198
- Buarremon brunneinucha*, 289–301
- B. torquatus*, 289–301
- Bubo bubo*, 35–45
- B. scandiaca*, 35–45
- B. virginianus*, 950–962, 1081–1093
- Bubulcus coromanda*, 985–995
- B. coromandus*, 985–995
- B. ibis*, 192–202, 985–995
- B. i. coromandus*, 985–995
- Bucconidae, 88–102
- Bucephala clangula*, 148–155
- buffer strips, 726–737
- Bullfinch, Eurasian, 372–379
- Bulweria bulwerii*, 985–995
- Bunting, Corn, 1226–1237; Indigo, 726–737, 865–876, 1040–1055, 1103–1121; Lark, 404–414, 1081–1093; Lazuli, 877–893; Yellow-throated, 985–995
- Buphagus erythrorhynchus*, 1155–1160
- Burden, Christina M., see Henson, Shandelle M., et al.
- Burnett, J. Alexander, A Passion for Wildlife: The History of the Canadian Wildlife Service (rev.), 273–275
- Burt, Edward H., Jr., see Goldstein, Gerald, et al.
- Buteo jamaicensis*, 738–751, 950–962
- B. lineatus*, 738–751
- B. platypterus*, 88–102
- B. rufogscus*, 1226–1237
- Buteogallus urubitinga*, 88–102
- Butorides striata*, 985–995
- B. striatus*, 985–995
- Buzzard, Jackal, 1226–1237
- Byers, Bruce E., et al., Extrapair paternity increases variability in male reproductive success in the Chestnut-sided Warbler (*Dendroica pensylvanica*), a socially monogamous songbird, 788–795
- Cacicus cela*, 88–102
- C. haemorrhous*, 88–102
- cacique, 224–229
- Calamospiza melanocorys*, 404–414, 1081–1093
- Calcarius lapponicus*, 435–445, 1238–1249
- C. ornatus*, 1081–1093, 1130–1145
- C. pictus*, 446–451
- Calidris alba*, 333–344
- C. alpina pacifica*, 170–177, 333–344
- C. canutus*, 1094–1102

- C. c. islandica*, 333-344
C. fuscicollis, 333-344, 777-787
C. maritima, 752-766
C. melanotos, 777-787
C. pusilla, 333-344
Callipepla squamata, 1081-1093
Calonectris diomedea, 847-864
C. leucomelas, 847-864
Campephilus rubricollis, 88-102
Campylopterus largipennis, 88-102
Campylorhamphus procurviroides, 88-102
Campylorhynchus, 1199-1207
C. brunneicapillus, 1081-1093
C. zonatus, 435-445
 Canary, 184-191
 canopy avifauna, 88-102
 Canterbury, Ronald A., see Shapiro, Leo H., et al.
 Canvasback, 170-177, 333-344
Capito niger, 88-102
 Capitonidae, 88-102
Caprimulgus carolinensis, 738-751
C. indicus, 985-995
 capture, 354-360
 capture-mark-recapture models, 15-22
 Cardinal, Northern, 74-87, 726-737, 738-751, 865-876, 1250-1261
 Cardinalinae, 88-102
Cardinalis cardinalis, 74-87, 435-445, 726-737, 738-751, 865-876, 1250-1261
C. sinuatus, 1081-1093
Carduelis carduelis, 1155-1160
C. chloris, 372-379, 473-485, 1155-1160
C. lawrencei, 824-836
C. pinus, 435-445
C. spinus, 1155-1160
C. tristis, 1250-1261
Carpodacus erythrinus, 361-371
C. mexicanus, 652-655
Caryothraustes canadensis, 88-102
 Catbird, Gray, 74-87, 726-737, 738-751, 865-876, 1040-1055, 1094-1102, 1250-1261
 Catharacta, 243-252
C. antarctica, 243-252
C. chilensis, 243-252
Cathartes melambrotus, 88-102
 Cathartidae, 88-102
Catharus, 88-102, 1040-1055
C. bicknelli, 446-451, 985-995
C. fuscescens, 1040-1055
C. guttatus, 46-57, 162-169, 738-751, 1094-1102
C. minimus, 985-995, 1040-1055
C. ustulatus, 162-169, 1040-1055
 Catry, Paulo, et al., Sustained fast travel by a Gray-headed Albatross (*Thalassarche chrysostoma*) riding an Antarctic storm, 1208-1213
Celeus torquatus, 88-102
C. undatus, 88-102
 censusing, 541-547
Cephalopterus ornatus, 1155-1160
Cephus, 148-155
C. grylle, 35-45, 463-472
Cercomacra tyrannina, 486-496
Cerorhinca monocerata, 452-462, 463-472
Certhia, 1155-1160
C. americana, 435-445
 Certhiidae, 1155-1160
 Cerulean Warbler Technical Group, 7-14
Ceryle, 985-995
C. alcyon, 35-45
C. guttulata, 985-995
C. lugubris, 985-995
C. rudis, 35-45
C. torquata, 985-995
C. torquatus, 985-995
 Cerylinae, 35-45
Cettia diphone, 1238-1249
 Chachalaca, Chaco, 682-694; Dusky-legged, 682-694; Sickie-winged, 682-694; Wattled, 682-694
Chaetura brachyura, 88-102
C. chapmani, 88-102
C. spinicauda, 88-102
 Chaffinch, Common, 1238-1249
Chamaepetes goudotii, 682-694
 Charadriiformes, 354-360
Charadrius leschenaultii, 985-995
C. mongolus, 985-995
C. montanus, 526-540
 Chat, Yellow-breasted, 526-540
Chelodactylus tenebrosus, 88-102
 Chen, 777-787
C. caerulescens, 587-599
C. c. atlantica, 777-787, 917-929
C. c. caerulescens, 333-344, 917-929
C. canagica, 985-995
 Chesser, R. Terry, Systematics, evolution, and biogeography of the South American ovenbird genus *Cinclodes*, 752-766
 Chickadee, Black-capped, 1040-1055, 1070-1080; Carolina, 74-87, 738-751, 865-876; Mountain, 777-787
 chicken, 308-317, 318-332, 372-379, 587-599, 656-659, 660-681, 717-725, 777-787, 847-864, 930-949, 1187-1198
 Chiffchaff, 110-117
 Chilton, Glen, review by, 982-983
 Chipley, Robert M., et al., The American Bird Conservancy Guide to the 500 Most Important Bird Areas in the United States (rev.), 977-978
Chiroxiphia caudata, 1155-1160
Chlorocichla falckensteini, 1155-1160
Chlorophanes spiza, 88-102
Chondestes grammacus, 1081-1093
Chordeiles acutipennis, 1081-1093
C. minor, 1081-1093
Chrysococcyx caprius, 1172-1186
 Chuck-will's-widow, 738-751

- Cicero, Carla, In Memoriam: Ned Keith Johnson, 1932–2003, 600–602
- Cicero, Carla, see Banks, Richard C., et al.
- Cinclidae, 1155–1160
- Cinclodes, 752–766
- C. antarcticus*, 752–766
- C. atacamensis*, 752–766
- C. comechingonus*, 752–766
- C. excelsior*, 752–766
- C. fuscus*, 752–766
- C. f. riojana*, 752–766
- C. nigrofumosus*, 752–766
- C. olrogi*, 752–766
- C. oustaleti*, 752–766
- C. pabsti*, 752–766
- C. palliatus*, 752–766
- C. patagonicus*, 752–766
- C. taczanowskii*, 752–766
- Cinclodes, Bar-winged, 752–766; Blackish, 752–766; Cordoba, 752–766; Dark-bellied, 752–766; Gray-flanked, 752–766; Long-tailed, 752–766; Olrog's, 752–766; Royal, 752–766; Seaside, 752–766; Stout-billed, 752–766; Surf, 752–766; White-bellied, 752–766; White-winged, 752–766
- Cinclus cinclus*, 1155–1160
- Circus aeruginosus*, 361–371
- Cisticola galactotes*, 1155–1160
- Cisticolidae, 1155–1160
- Cistothorus palustris*, 391–403, 1226–1237
- citizen scientists, 118–133
- cladistic analyses, 422–434
- Clark, Robert G., see Drever, Mark C., et al.
- classification, 985–995
- Climacteridae, 1155–1160
- Climacteris leucophaea*, 1155–1160
- climate variation, 497–508
- climatic variability, 1214–1225
- clustered breeding, 877–893
- clutch, — enlargement, 557–564; — initiation, 345–353; — size, 103–109, 289–301, 557–564, 587–599, 695–705, 1070–1080
- clutch-size evolution, 289–301
- coastal sage-scrub habitat, 904–916
- Coccythraustes coccythraustes*, 1155–1160
- Coccyzus americanus*, 1040–1055
- C. erythrophthalmus*, 1040–1055
- C. melacoryphus*, 985–995
- Cock-of-the-Rock, 391–403
- Coereba flaveola*, 88–102, 652–655
- Coerebinae, 88–102
- coevolution, parasite–host, 1172–1186
- coevolutionary cycles, 1172–1186
- Cohen, Emily B., and Lindell, Catherine A., Survival, habitat use, and movements of fledgling White-throated Robins (*Turdus assimilis*) in a Costa Rican agricultural landscape, 404–414
- Colaptes auratus*, 509–514, 1081–1093
- Collar, Nigel J., Threatened Birds of Asia: The BirdLife International Red Data Book (rev.), 619–622
- colonial dynamics, 192–202
- coloniality, 257–262, 811–823, 1122–1129
- colonies, heron, 192–202
- colony, — site selection, 192–202; — size, 1122–1129
- Colorhamphus parvirostris*, 515–525
- Columba araucana*, 515–525
- C. livia*, 103–109, 837–846
- C. plumbea*, 88–102
- C. subvinacea*, 88–102
- Columbidae, 88–102
- commodity selection hypothesis, 257–262
- communication, 184–191
- competition, direct, 806–810
- Conopias albivittata*, 985–995
- C. albivittatus*, 985–995
- C. parva*, 88–102, 985–995
- C. parvus*, 985–995
- Conquest, Loveday L., see Hamel, Nathalie J., et al.
- conservation, 7–14, 15–22, 23–34, 58–67, 88–102, 118–133, 192–202, 243–252, 380–390, 515–525, 548–556, 717–725, 806–810, 917–929, 1103–1121, 1130–1145
- Contopus borealis*, 88–102
- C. caribaeus*, 985–995
- C. sordidulus*, 824–836
- C. virens*, 865–876, 1040–1055
- Contreras, Thomas A., see Sieving, Kathryn E., et al.
- convergent evolution, 548–556
- Cooke, Fred, review by, 1301
- Cooper, Robert J., review by, 622–623, 977–978
- cooperative breeding, 811–823, 1199–1207
- coots, 1122–1129
- Copsychus saularis*, 435–445, 985–995
- copulation-solicitation displays, 824–836
- Coraciiformes, 811–823
- Coragyps atratus*, 88–102
- Corapipo gutturalis*, 88–102
- Corcorax melanorhamphos*, 1155–1160
- Cordon-bleu, Blue-capped, 156–161
- Cormorant, Imperial, 243–252; Neotropic, 243–252
- cormorants, 35–45, 148–155
- corticosterone, 276–277, 1094–1102, 1238–1249
- Corvidae, 162–169, 361–371, 1155–1160
- corvids, 361–371
- Corvus brachyrhynchos*, 170–177, 354–360, 435–445, 865–876, 950–962
- C. caurinus*, 435–445
- C. corax*, 162–169, 361–371, 637–651
- C. corone*, 361–371, 372–379, 452–462
- C. cryptoleucus*, 1081–1093
- C. frugilegus*, 35–45
- C. macrorhynchos*, 452–462
- C. monedula*, 372–379, 1155–1160
- C. moneduloides*, 509–514
- Corydon sumatranus*, 1155–1160
- Cosmopsarus regius*, 1155–1160

- Cossypha*, 23–34
C. cyanocampter, 435–445
Cotinga cayana, 88–102
C. cotinga, 88–102
cotingas, 88–102
Cotingidae, 88–102
Coturnix japonica, 170–177, 333–344
 Coues Award, 2003: Donald E. Kroodsma, 263–264
 courtship song, 824–836
 Covas, Rita, et al., Juvenile and adult survival in the Sociable Weaver (*Philetairus socius*), a southern-temperate colonial cooperative breeder in Africa, 1199–1207
 Cowbird, 767–776; Bronzed, 1172–1186; Brown-headed, 7–14, 308–317, 354–360, 726–737, 767–776, 824–836, 865–876, 877–893, 1081–1093, 1103–1121, 1130–1145, 1172–1186; Giant, 1172–1186; Shiny, 1172–1186
Cracidae, 88–102, 230–242, 682–694
 cranes, 1–6
Crax, 682–694
C. alberti, 682–694
C. alector, 682–694
C. blumenbachii, 682–694
C. daubentoni, 682–694
C. fasciolata, 682–694
C. globulosa, 682–694
C. rubra, 682–694
Creatophora cinerea, 435–445
 Creeper, Hawaii, 548–556
Criniger, 23–34
 cross-fostering, 1172–1186
 Crow, American, 170–177, 354–360, 865–876, 950–962; Carrion, 361–371, 372–379, 452–462; Jungle, 452–462; New Caledonian, 509–514
 Crowe, Timothy M., see Bowie, Rauri C. K., et al.
 Croxall, John P., see Catry, Paulo, et al.
 cryptic species, 415–421
Cryptospiza, 23–34
 cuckoldry, 788–795
 Cuckoo, 1155–1160; Black-billed, 1040–1055; Common, 1172–1186; Diederick, 1172–1186; Yellow-billed, 1040–1055
Cuculidae, 88–102, 1155–1160
Cuculus canorus, 1172–1186
 Cuisin, Jacques, see Villard, Pascal, and —
 Currasow, 682–694; Alagoas, 682–694; Bare-faced, 682–694; Black, 682–694; Blue-knobbed, 682–694; Crestless, 682–694; Great, 682–694; Helmeted, 682–694; Horned, 682–694; Nocturnal, 682–694; Razor-billed, 682–694; Red-billed, 682–694; Salvin's, 682–694; Wattled, 682–694; Yellow-billed, 682–694
Cyanerpes caeruleus, 88–102
C. cyaneus, 88–102
C. nitidus, 88–102
Cyanicterus cyanicterus, 88–102
Cyanistes caeruleus, 345–353
Cyanocitta cristata, 354–360, 435–445, 738–751, 865–876, 877–893
C. stelleri, 435–445
Cyanocorax morio, 435–445
Cyanopsitta spixii, 230–242
Cyclarhis gujanensis, 88–102
Cymbilaimus lineatus, 88–102
Cymbirhynchus macrorhynchus, 1155–1160
 dabbling ducks, 497–508, 695–706, 1161–1171
Dacnis cayana, 88–102
D. lineata, 88–102
 Dadsell, Ted, The Selborne Pioneer: Gilbert White as Naturalist and Scientist, a Re-examination (rev.), 270–272
Daption capense, 422–434
Daptrius americanus, 88–102
D. ater, 88–102
 darters, 35–45
 Darwin's finches, 660–681
 Davis, Lloyd S., and Renner, Martin, Penguins (rev.), 1293–1296
 Davis, Stephen K., Area sensitivity in grassland passerines: Effects of patch size, patch shape, and vegetation structure on bird abundance and occurrence in southern Saskatchewan, 1130–1145
 dawn chorus, 184–191
 Dawson, Alistair, and Newton, Ian, Use and validation of a molt score index corrected for primary-feather mass, 372–379
 Dawson, Deanna K., see Hamel, Paul B., et al.
 Dean, W. Richard J., In Memoriam: Richard Liversidge, 1926–2003, 603
 DeAngelis, Donald L., see Dreitz, Victoria J., et al.
 Deguchi, Tomohiro, et al., Proximate factors determining age and mass at fledging in Rhinoceros Auklets (*Cerorhinca monocerata*): Intra- and interyear variations, 452–462
Delichon urbica, 985–995, 1155–1160, 1187–1198
D. urbicum, 985–995
 Delpont, Wayne, et al., Structure of an African Red-billed Hornbill (*Tockus erythrorhynchus rufirostris* and *T. e. damarensis*) hybrid zone as revealed by morphology, behavior, and breeding biology, 565–586
 demography, 118–133, 1056–1069, 1103–1121
Dendragapus canadensis, 162–169
Dendrexetastes rufigula, 88–102
Dendrocincla cf. fuliginosa, 1155–1160
Dendrocolaptidae, 88–102, 752–766
Dendroica, 15–22, 224–229, 610–618
D. aestiva, 610–618
D. auduboni, 610–618
D. caerulescens, 15–22, 302–307, 473–485, 788–795, 1040–1055, 1094–1102
D. castanea, 162–169, 877–893, 1040–1055
D. cerulea, 7–14, 15–22

- D. coronata*, 15–22, 46–57, 74–87, 162–169, 289–301, 610–618, 738–751, 1040–1055, 1094–1102
D. discolor, 738–751, 1040–1055, 1250–1261
D. dominica, 610–618, 1040–1055
D. fusca, 1040–1055
D. kirtlandii, 877–893
D. magnolia, 162–169, 610–618, 1040–1055
D. nigrescens, 610–618
D. occidentalis, 610–618, 1014–1018
D. palmarum, 738–751
D. pensylvanica, 788–795, 1040–1055
D. petechia, 788–795, 985–995, 1238–1249
D. p. aestiva, 1238–1249
D. p. parkesi, 1238–1249
D. pinus, 71–87, 738–751
D. striata, 88–102, 162–169, 435–445, 1040–1055
D. tigrina, 1040–1055
D. townsendi, 15–22, 610–618, 1014–1018
D. virens, 162–169, 610–618, 1040–1055
 density-dependent effects, 210–223
Derophtus accipitrinus, 88–102, 230–242
 desert grasslands, 1081–1093
 Desrochers, André, see Ibarzabal, Jacques, and —
 detectability, 865–876
 detection probabilities, 865–876
 development, 1187–1198
 developmental conversion, 1187–1198
 Diaz, Iván A., see Reid, Sharon, et al.
Dicaeum hirundinaceum, 1155–1160
 Dickcissel, 435–445, 767–776, 1250–1261
Dicrurus adsimilis, 1155–1160
D. paradiseus, 435–445
 diet, 170–177, 333–344, 509–514, 695–705
 diet-tissue fractionation, 170–177
Diglossa lafresnayii, 1155–1160
Diomedea, 1–6
D. chlororhynchos, 463–472
D. exulans, 1208–1213
D. melanophris, 422–434, 463–472
Diomedidae, 1–6
Diopsittaca, 230–242
D. nobilis, 230–242
Discosura longicauda, 88–102
 disease, 717–725
 dispersal, 682–694, 1146–1154; delayed, 1199–1207;
 — distances, 1103–1121; male, 930–949; natal,
 404–414, 894–903; postfledging, 203–209
 diurnal abundance dynamics, 380–390
 diversification, 224–229, 230–242
Dives atrovioleacea, 985–995
D. atrovioleaceus, 985–995
 DNA sequencing, 230–242
 Dobkin, David S., see George, T. Luke, and —
Dolichonyx oryzivorus, 767–776, 1103–1121, 1250–1261
 dominance rank, 1070–1080
 Dorst, Jean, 1924–2001, In Memoriam, 1289–1290
 double-brooding, 796–805
 double-observer modeling, 865–876
 Dove, Carla J., et al., In Memoriam: Roxie Collie
 Laybourne, 1910–2003, 1282–1285
 Dove, Mourning, 354–360, 1081–1093, 1172–1186;
 Rock, 837–846; White-winged, 1081–1093
 Dreitz, Victoria J., et al., Effects of natal departure
 and water level on survival of juvenile Snail
 Kites (*Rostrhamus sociabilis*) in Florida, 894–903
Drepanidae, 548–556
 Drever, Mark C., et al., Decline of duck nest success
 revisited: Relationships with predators and
 wetlands in dynamic prairie environments,
 497–508
Dryobates, 985–995
Dryocopus lineatus, 88–102
D. pileatus, 509–514
 du Plessis, Morné A., see Radford, Andrew N., and —
 Duck, American Black, 930–949; Eastern Spot-billed,
 930–949; Hawaiian, 930–949; Mexican, 930–949;
 Mottled, 930–949; Pacific Black, 930–949;
 Phillipine, 930–949; Spot-billed, 930–949; Wood,
 148–155
Dumetella carolinensis, 74–87, 435–445, 726–737, 738–
 751, 865–876, 1040–1055, 1094–1102, 1155–1160,
 1250–1261
 Dunlin, 170–177, 333–344
 Dunn, Jon L., see Banks, Richard C., et al.
 Dunnock, 446–451
 Durant, Joël M., et al., More eggs the better: Egg
 formation in captive Barn Owls (*Tyto alba*),
 103–109
 Eagle, 88–102; Bald, 463–472; Tawny, 1081–1093
 Eberhard, Jessica R., and Bermingham, Eldredge,
 Phylogeny and biogeography of the *Amazona*
 ochrocephala (Aves: Psittacidae) complex, 318–332
 Eckert, Cameron D., see Sinclair, Pamela H., et al.
 ecological, — constraints, 1238–1249; — dynamics,
 380–390; — traps, 162–169
 edge, — avoidance, 767–776; — effect, 361–371, 726–
 737; — habitats, 904–916
 Eens, Marcel, see Gorissen, Leen, and —
 egg, — breakage, 148–155; — cooling rates, 777–787;
 — formation, 103–109; — laying, 557–564;
 — period, 148–155; — production, 695–705;
 — rejection, 1172–1186; — temperature, 587–
 599, 777–787; — tossing, 811–823; — transfer,
 1122–1129
 eggshell, 148–155; — mass, 148–155; — thickness,
 148–155
 egg-weight, relative, 308–317
 Egret, Cattle, 192–202, 985–995; Little, 192–202
Egretta garzetta, 192–202
 Eider, Common, 333–344, 777–787
 El Niño event, 904–916
Elaenia albiceps, 515–525
E. chiriquensis, 1005–1013
E. flavogaster, 435–445
E. strepera, 1005–1013

- Elaenia, Lesser, 1005–1013; Slaty, 1005–1013; White-crested, 515–525
Elanoides forficatus, 88–102, 404–414
Elanus leucurus, 904–916
Emberiza citrinella, 1155–1160
E. elegans, 985–995
 Emberizidae, 88–102, 1081–1093
 emberizids, 289–301
 emigration, 930–949
 Emlen, Stephen T., and Wrege, Peter H., Size dimorphism, intrasexual competition, and sexual selection in Wattled Jacana (*Jacana jacana*), a sex-role-reversed shorebird in Panama, 391–403
Empidonax alnorum, 1040–1055
E. flaviventris, 1040–1055
E. minimus, 877–893, 1040–1055
E. traillii, 656–659, 1040–1055, 1103–1121
E. virescens, 707–716, 726–737, 865–876
Empidonax aurantioatrocristatus, 1005–1013
E. varius, 88–102
 endemic, 23–34, 1031–1039
 energetic condition, 1094–1102
 energetics, 1187–1198
 energy, — budgets, 1208–1213; — reserves, 1094–1102
 energy-delivery rates, 1161–1171
 environmental, — conditions, 1094–1102; — factors, 894–903, 1199–1207; — variability, 257–262
Eopsaltria australis, 1155–1160
Eremophila alpestris, 435–445, 1081–1093, 1130–1145, 1155–1160
Erithacus, 23–34
E. rubecula, 1155–1160
 Erratum, 276–277
Erythropgia, 23–34
 escape cover, 515–525
 estimating nest survival, 134–147
Estrilda melpoda, 435–445
 estrildid finches, 156–161, 435–445
 Estrildidae, 435–445
Eudypetes, 148–155, 422–434
E. chrysocome chrysocome, 422–434
E. c. moseleyi, 422–434
E. chrysolophus, 148–155, 422–434
E. pachyrrhynchus, 422–434
E. robustus, 422–434
E. schlegeli, 422–434
E. sclateri, 422–434
Eudypula, 422–434
E. minor, 422–434
E. m. albosignata, 422–434
E. m. minor, 422–434
Eugralla paradoxa, 515–525
 Eupasseridae, 1155–1160
Euphagus carolinus, 162–169
E. cyanocephalus, 435–445
E. cayennensis, 88–102
E. chrysopasta, 88–102
E. hirundinacea, 435–445
E. minuta, 88–102
Euplectes, 1226–1237
E. ardens, 1226–1237
E. jacksoni, 1226–1237
E. orix, 435–445, 1226–1237
 European passeriform avifauna, 1155–1160
 Eurylaimidae, 1155–1160
Eurynorhynchus pygmeus, 985–995
 Evans, Monica Ann, In Memoriam: H. Lewis Batts, Jr., 1922–2001, 970
 Evans Ogden, Lesley J., et al., Blood isotopic ($\delta^{13}\text{C}$ and $\delta^{15}\text{N}$) turnover and diet-tissue fractionation factors in captive Dunlin (*Calidris alpina pacifica*), 170–177
 evolution, 752–766; — of coloniality, 257–262; — of migration, 610–618
 evolutionary, — biology, 637–651; — variation, 435–445
 exposure days, 707–716
 extrapair, — copulations, 877–893; — fertilization, 302–307, 788–795; — paternity, 302–307, 473–485, 788–795
 fairy-bluebirds, 435–445
 Fairy-wren, White-winged, 652–655
Falco columbarius, 1081–1093
F. femoralis, 1081–1093
F. f. septentrionalis, 1081–1093
F. mexicanus, 1081–1093
F. peregrinus, 148–155, 230–242, 1081–1093
F. rufigularis, 88–102
F. sparverius, 738–751, 837–846, 904–916, 1081–1093, 1187–1198
F. timunculus, 103–109
 Falcon, 35–45, 88–102; Aplomado, 1081–1093; Northern Aplomado, 1081–1093; Peregrine, 148–155, 1081–1093; Prairie, 1081–1093
 Falconidae, 53–45, 88–102
Falcula palliata, 1155–1160
 fall stopover, 1040–1055
 fast travel, 1208–1213
 fatty acids, 435–445
 feather, — coloration, 652–655; — mass, 372–379; — parasites, 1262–1268; — waxes, 435–445
 feathers, 372–379, 652–655, 656–659; flight, 35–45
 fecundity, 788–795, 1056–1069, 1214–1225; annual, 796–805, 1250–1261
 Fedy, Bradley C., and Stutchbury, Bridget J. M., Territory switching and floating in White-bellied Antbird (*Myrmeciza longipes*), a resident tropical passerine in Panama, 486–496
 feeding, — apparatus, 509–514; — rates, 345–353
 female, — call, 184–191; — nutrition, 695–705
 Fenwick, George H., see Chipley, Robert M., et al.
 Ferguson, J. Willem H., see Delpont, Wayne, et al.
Ficedula albicollis, 110–117, 717–725
F. hypoleuca, 345–353, 837–846, 1155–1160, 1226–1237, 1238–1249

- F. parva*, 637–651
 fidelity, natal-site, 930–949; site, 354–360, 1103–1121;
 territory, 308–317, 1103–1121
 Finch, House, 652–655; Melba, 156–161; Zebra,
 473–485
 fire frequency, 74–87
 Firecrown, Green-backed, 515–525
 Firefinch, 156–161; Red-billed, 156–161
 fitness, 257–262
 Fjeldså, Jon, see Beresford, Pamela, et al.
 Fjeldså, Jon, see Bowie, Rauri C. K., et al.
 Flamingo, Greater, 541–547
 fledging, — age, 452–462, 463–472; — mass, 452–462;
 — success, 557–564
 Fleischer, Robert C., see Shapiro, Leo H., et al.
 Flicker, Northern, 509–514, 1081–1093
 flight, — ability, 345–353; — costs, 1208–1213;
 — performance, 345–353; — whistles, 824–836
 floaters, 486–496
Florisuga mellivora, 88–102
 Flory, Kelly R., see Goldstein, Gerald, et al.
 flushing, 354–360
 Flycatcher, 88–102, 1005–1013; Acadian, 707–716, 726–
 737, 865–876; Alder, 1040–1055; Ash-throated,
 1081–1093; Collared, 110–117, 717–725; Crowned
 Slaty, 1005–1013; Dark-sided, 985–995; Gray-
 spotted, 985–995; Gray-streaked, 985–995; Great
 Crested, 738–751, 1040–1055; Least, 877–893,
 1040–1055; Pied, 345–353, 837–846, 1226–1237,
 1238–1249; Red-breasted, 637–651; Siberian,
 985–995; Spotted, 985–995; Swainson's, 610–618,
 1005–1013; Vermilion, 1005–1013, 1081–1093;
 Willow, 656–659, 1040–1055, 1103–1121; Yellow-
 bellied, 1040–1055
 follicles, 103–109
 food, — abundance, 210–223, 557–564; — availability,
 74–87, 463–472, 515–525, 1199–1207; — limitation,
 289–301, 446–451; — resource, 515–525
 forage availability, 210–223
 foraging, 509–514, 548–556; — competition, 178–183;
 — performance, 345–353; — trip, 1208–1213
 forest, — canopy, 88–102; — edges, 162–169;
 — habitats, 1040–1055; managed, 162–169;
 — robins, 23–34
 forest-boundary crossing, 738–751
 formic acid, 1262–1268
 Formicariidae, 88–102, 515–525, 1155–1160
Formicarius, 435–445, 1155–1160
F. a. alis, 435–445
Forpus passerinus, 717–725
 Forrester, Donald J., and Spalding, Marilyn G.,
 Parasites and Diseases of Wild Birds in Florida
 (rev.), 272–273
 Forsman, Eric D., see Kelly, Elizabeth G., and —
 Fort, Kevin T., and Otter, Ken A., Effects of habitat
 disturbance on reproduction in Black-capped
 Chickadees (*Parus atricapillus*) in northern
 British Columbia, 1070–1080
 fossil record, 1155–1160
 fragmented landscapes, 738–751
Fratercula arctica, 35–45, 452–462, 463–472
F. cirrhata, 35–45, 463–472
F. corniculata, 35–45
Fringilla, 435–445
F. coelebs, 435–445, 1155–1160, 1238–1249
F. montifringilla, 1155–1160, 1238–1249
 Fringillidae, 1155–1160
 fringillids, 435–445
 Frogmouth, Marbled, 541–547
 frugivores, 88–102, 1040–1055
 fruit, — abundance, 46–57, 74–87; — availability,
 1040–1055; — pulp biomass, 46–57;
 — resources, 74–87
 Fullagar, Peter, In Memoriam: Stephen Marchant,
 1912–2003, 971
 Fulmar, Northern, 289–301
Fulmarus glacialis, 289–301
 fungicidal activity, 1262–1268
 Furnariidae, 88–102, 515–525, 752–766, 1155–1160
 furnariids, 752–766
Furnarius rufus, 752–766, 1155–1160
 Gadwall, 497–508, 587–599, 695–706, 950–962
Galbula dea, 88–102
 Galbulidae, 88–102
 Gale, George A., review by, 1299–1300
Galerida cristata, 1155–1160
 Galliformes, 23–34, 230–242, 308–317, 682–694
 Galloanserae, 230–242
Gallus, 1–6
G. gallus, 230–242, 308–317, 318–332, 372–379,
 660–681, 717–725, 777–787, 847–864, 930–949,
 1187–1198
G. g. domesticus, 656–659
 Galusha, Joseph G., see Henson, Shandelle M., et al.
 García Borboroglu, Pablo, and Yorio, Pablo, Habitat
 requirements and selection by Kelp Gulls (*Larus*
dominicanus) in central and northern Patagonia,
 Argentina, 243–252
Garrulax leucolophus, 435–445, 1155–1160
Garrulus glandarius, 35–45, 361–371
 gas chromatography-mass spectrometry, 435–445
Gavia, 422–434
G. stellata, 422–434
 Gaviidae, 35–45
 Gaviiformes, 422–434
 Gavin, Thomas A., see Bollinger, Eric K., and —
 gene flow, sex-biased, 930–949
 genetic, — divergence, 660–681; — variability,
 302–307
Geococcyx californianus, 1081–1093
 George, T. Luke, and Dobkin, David S., Effects of
 Habitat Fragmentation on Birds in Western
 Landscapes: Contrasts with Paradigms from the
 Eastern United States (rev.), 978–980
Geositta, 752–766

- G. cunicularia*, 752–766
Geospiza, 660–681
Geothlypis aequinoctialis, 985–995
G. trichas, 435–445, 738–751, 788–795, 865–876, 1040–1055
Geotrygon caniceps, 985–995
G. leucometopia, 985–995
 Giannini, Norberto P., and Bertelli, Sara, Phylogeny of extant penguins based on integumentary and breeding characters, 422–434
 Gill, Frank B., Blue-winged Warblers (*Vermivora pinus*) versus Golden-winged Warblers (*V. chrysoptera*), 1014–1018
 Gjerdrum, Carina, Parental provisioning and nestling departure decisions: A supplementary feeding experiment in Tufted Puffins (*Fratercula cirrhata*) on Triangle Island, British Columbia, 463–472
Glaucidium hardyi, 88–102
G. nanum, 515–525
G. passerinum, 541–547
Glaucis aenea, 985–995
G. aeneus, 985–995
G. hirsuta, 985–995
G. hirsutus, 985–995
 Gnatcatcher, Black-tailed, 1081–1093; Blue-gray, 738–751, 1040–1055
 Godwit, Bar-tailed, 1094–1102
 Goethe, Friedrich Walter, 1911–2003, In Memoriam, 1292
 Goldbreast, 156–161
 Goldcrest, 877–893
 Goldeneye, Common, 148–155
 Goldfinch, American, 1250–1261; Lawrence's, 824–836
 Goldstein, Gerald, et al., Bacterial degradation of black and white feathers, 656–659
 Goose, 1–6, 68–73, 210–223, 777–787; Barnacle, 68–73; Cackling, 985–995; Canada, 985–995; Greater Snow, 777–787, 917–929; Lesser Snow, 333–344, 587–599, 652–655, 917–929
 Gorissen, Leen, and Eens, Marcel, Interactive communication between male and female Great Tits (*Parus major*) during the dawn chorus, 184–191
 Goshawk, African, 1226–1237
 gosling growth, 210–223
 Göth, Ann, and Vogel, Uwe, Is monogamy in the Polynesian Megapode (*Megapodius pritchardii*) related to its high relative egg-weight?, 308–317
 Grackle, 224–229, 1172–1186; Boat-tailed, 1172–1186; Carib, 1172–1186; Common, 1172–1186; Greater Antillean, 1172–1186; Great-tailed, 1081–1093, 1172–1186
Gracula religiosa, 435–445, 985–995, 1155–1160
 Graham, Frank, Jr., In Memoriam: Richard H. Pough, 1904–2003, 969
Grallina cyanoleuca, 1155–1160
 grassland birds, 767–776, 796–805, 1081–1093, 1130–1145
 Graves, Gary R., Testicular volume and asymmetry are age-dependent in Black-throated Blue Warblers (*Dendroica caerulescens*), 473–485
 grazing, — lawns, 210–223; — pressure, 210–223
 grebes, 35–45, 777–787
 Green, Jordan, see Perkins, Kelly A., et al.
 Greenberg, Cathryn H., see Borgmann, Kathi L., et al.
 Greenberg, Cathryn H., see Kwit, Charles, et al.
 Greenfinch, European, 372–379, 473–485
 greenlets, 88–102
 Greiner, Ellis C., see Smith, Ruth B., et al.
 Grosbeak, Black-headed, 1081–1093; Blue, 1081–1093; Rose-breasted, 1040–1055
 Ground-Tyrant, Cinnamon-bellied, 1005–1013
 Grouse, Spruce, 162–169
 growth, delayed, 1187–1198; early, 68–73; nestling, 463–472; — rates, 15–22, 452–462
 guans, 88–102, 682–694
Guarouba guarouba, 230–242
 Guillemot, 148–155; Black, 35–45, 463–472
 Gull, 243–252; Black-headed, 361–371; California, 950–962; Dolphin, 243–252; Glaucous-winged, 380–390; Kelp, 243–252; Lesser Black-backed, 243–252; Olog's, 243–252; Ring-billed, 170–177, 950–962; Slaty-backed, 452–462
 Gutiérrez, R. J., see LaHaye, William S., et al.
 Gymnogene, 1226–1237
Gymnostinops, 224–229
G. bifasciatus, 224–229
G. b. neivae, 224–229
G. b. yuracares, 224–229
G. guatemalensis, 224–229
G. montezuma, 224–229
 habitat, 1081–1093; — characteristics, 877–893; — disturbance, 1070–1080; — edges, 767–776, 950–962, 1250–1261; — fragmentation, 88–102, 361–371, 904–916, 1130–1145, 1214–1225; — heterogeneity, 877–893; — loss, 7–14, 15–22, 806–810; — mosaics, 738–751; — patch occupancy, 380–390; — patches, 767–776; — quality, 1040–1055, 1070–1080; — requirements, 243–252; — selection, 243–252, 1040–1055; — structure, 707–716; — use, 46–57, 74–87, 404–414, 1040–1055
 habitat-patch effects, 726–737
 Hackett, Shannon J., see Bowie, Rauri C. K., et al.
Haematoderus militaris, 88–102
 Hafner, Heinz, see Tourenq, Christophe, et al.
 Hahn, Thomas P., et al., Facultative altitudinal movements by Mountain White-crowned Sparrows (*Zonotrichia leucophrys oriantha*) in the Sierra Nevada, 1269–1281
Haliaeetus leucocephalus, 463–472
 Hamel, Nathalie J., et al., Effects of tagging on behavior, provisioning, and reproduction in the Common Murre (*Uria aalge*), a diving seabird, 1161–1171

- Hamel, Paul B., et al., How we can learn more about the Cerulean Warbler (*Dendroica cerulea*), 7–14
- Handrich, Yves, see Durant, Joël M., et al.
- Hansell, Mike H., see Hilton, Geoff M., et al.
- Harpagus bidentatus*, 88–102
- Harpya harpija*, 88–102
- Harrier, Eurasian Marsh, 361–371
- Harvey, Erica L., see Sweeney, Robert J., et al.
- hatching, — date, 452–462; — synchrony, 587–599
- Hawk, 35–45, 88–102, 877–893; Bicolored, 515–525; Cooper's, 738–751, 904–916; Red-shouldered, 738–751; Red-tailed, 738–751, 950–962; Sharp-shinned, 738–751, 837–846, 904–916
- Hayward, James L., see Henson, Shandelle M., et al.
- Hazler, Kirsten R., Mayfield logistic regression: A practical approach for analysis of nest survival, 707–716
- Heacker, Marcy, see Dove, Carla J., et al.
- health, 717–725
- Heinrich, Bernd, review by, 270–272
- Heliodoxa aurescens*, 973–976
- Heliothryx aurita*, 88–102, 985–995
- H. auritus*, 985–995
- H. barroti*, 985–995
- Helminthophaga lawrencei*, 1014–1018
- H. leucobronchialis*, 1014–1018
- Helmitheros vermivorus*, 985–995
- H. vermivorus*, 985–995, 1040–1055
- hematology, 717–725
- hematozoa, 837–846
- Hemignathus*, 1–6
- H. munroi*, 548–556
- H. virens*, 548–556
- Hemithraupis flavicollis*, 88–102
- Henson, Shandelle M., et al., Predicting dynamics of aggregate loafing behavior in Glaucous-winged Gulls (*Larus glaucescens*) at a Washington colony, 380–390
- herbivores, 210–223
- Heron, Squacco, 192–202
- Herpsilochmus*, 88–102
- H. dorsimaculatus*, 88–102
- Herzog, Mark P., and Sedinger, James S., Dynamics of foraging behavior associated with variation in habitat and forage availability in captive Black Brant (*Branta bernicla nigricans*) goslings in Alaska, 210–223
- Herzog, Mark P., see Sedinger, James S., et al.
- Herzog, Sebastian K., and Mazar Barnett, Juan, On the validity and confused identity of *Serpophaga griseiceps* Berlioz 1959 (Tyrannidae), 415–421
- Heteralocha acutirostris*, 178–183
- heterozygosity, 930–949
- heuristic parsimony analysis, 422–434
- Hill, Geoffrey E., A Red Bird in a Brown Bag: The Function and Evolution of Colorful Plumage in the House Finch (rev.), 267–269
- Hill, Geoffrey E., see Shawkey, Matthew D., and —
- Hilton, Geoff M., et al., Using artificial nests to test importance of nesting material and nest shelter for incubation energetics, 777–787
- Himatione sanguinea*, 435–445
- Hippolais icterina*, 1155–1160
- Hirundinidae, 88–102, 1155–1160, 1187–1198
- Hirundo ariel*, 302–307
- H. pyrrhonota*, 435–445
- H. rustica*, 1122–1129, 1155–1160, 1214–1225
- history of biology, 637–651
- Hobson, Keith A., see Evans Ogden, Lesley J., et al.
- Hobson, Keith A., see Morrison, R. I. Guy, and —
- Hoi, Christine, see Hoi, Herbert, et al.
- Hoi, Herbert, et al., Clutch enlargement in Lesser Gray Shrikes (*Lanius minor*) in Slovakia when food is superabundant: A maladaptive response?, 557–564
- Holberton, Rebecca L., see Long, Jennifer A., and Holberton, Rebecca L., see Wilson, C. Morgan, and —
- holotype, 1031–1039
- Holyoak, David T., Nightjars and Their Allies: The Caprimulgiformes (rev.), 622–623
- home range, 58–67, 446–451
- honeycreepers, 88–102, 548–556; Hawaiian, 1–6, 548–556
- Hornbill, African Red-billed, 565–586; Damaraland Red-billed, 565–586; Monteiro's, 565–586; Red-billed, 565–586
- host defenses, 1172–1186
- Houston, Mary, see Houston, Stuart, et al.
- Houston, Stuart, et al., Eighteenth-Century Naturalists of Hudson Bay (rev.), 1301
- Howerth, Elizabeth W., review by, 272–273
- Huet-huet, Black-throated, 515–525
- Hughes, Colin R., see Sheridan, Jennifer A., et al.
- Hughes, Nancy L., see Sinclair, Pamela H., et al.
- Huia, 178–183
- Hummingbird, 35–45, 88–102, 435–445, 973–976; Ruby-throated, 877–893, 1040–1055
- Hunt, W. Grainger, see Macias-Duarte, Alberto, et al.
- hybrid, 156–161; — zone, 565–586, 1014–1018
- hybridization, 660–681, 806–810, 1019–1030; asymmetric, 930–949
- Hydrophasianus chirurgus*, 391–403
- Hylocharis*, 985–995
- H. sapphirina*, 88–102
- Hylocichla mustelina*, 46–57, 354–360, 404–414, 726–737, 824–836, 865–876, 1040–1055
- Hyloctistes subulatus*, 985–995
- H. virgatus*, 985–995
- Hylophilus muscicapinus*, 88–102
- Hylophylax naevia*, 985–995
- H. naevioides*, 985–995
- H. naevius*, 985–995
- Hypargos niveoguttatus*, 156–161
- hypotarsus, 1155–1160
- Hypsipetes madagascariensis*, 1155–1160

- Ibarzabal, Jacques, and Desrochers, André, A nest predator's view of a managed forest: Gray Jay (*Perisoreus canadensis*) movement patterns in response to forest edges, 162-169
- Ichida, Jann M., see Goldstein, Gerald, et al.
- Icteria virens*, 435-445, 526-540
- Icterinae, 88-102, 1172-1186
- Icterini, 224-229
- Icterus chrysiocephalus*, 88-102
- I. dominicensis*, 1155-1160
- I. fortis*, 985-995
- I. galbula*, 224-229, 435-445, 1040-1055
- I. parisorum*, 1081-1093
- I. spurius*, 985-995
- Ictinia plumbea*, 88-102
- identification, behavioral, 156-161; genetic, 156-161
- immune system, 837-846
- immunoglobulin-G levels, 717-725
- imprinting, 156-161
- incubation, 557-564, 587-599, 777-787; — energetics, 777-787
- Indigobird, 156-161; Purple, 156-161; Twinspot, 156-161; Village, 156-161
- individual, — quality, 345-353; — recognition, 541-547
- information-theoretic, — approach, 726-737; — techniques, 526-540
- insectivore-frugivores, 88-102
- insectivores, 88-102, 1187-1198
- integumentary characters, 422-434
- intersexual communication, 184-191
- intraclutch egg-variation, 1172-1186
- intrasexual competition, 391-403
- introgression, 565-586, 930-949, 1014-1018; asymmetric mitochondrial, 1019-1030; reciprocal, 1019-1030
- invertebrate abundance, 515-525
- investigator disturbance, 354-360
- Irediparra gallinacea*, 391-403
- Irena*, 435-445
- I. puella*, 435-445, 1155-1160
- Irenidae, 1155-1160
- isolating mechanisms, 806-810
- isotope models, 170-177
- isotopic turnover, 170-177
- Ixoreus*, 435-445
- I. naevius*, 435-445
- jacamars, 88-102
- Jacana jacana*, 391-403
- J. spinosa*, 391-403
- Jacana, African, 391-403; Bronze-winged, 391-403; Comb-crested, 391-403; Lesser, 391-403; Northern, 391-403; Pheasant-tailed, 391-403; Wattled, 391-403
- Jacaniidae, 391-403
- Jackdaw, Eurasian, 372-379
- Jaeger, Parasitic, 652-655
- Jahn, Alex E., et al., Reflections across hemispheres: A system-wide approach to New World bird migration, 1005-1013
- James, Frances C., and Mezey, Jason G., review by, 269-270
- Jay, Blue, 354-360, 738-751, 865-876, 877-893; Eurasian, 35-45, 361-371; Gray, 162-169; Siberian, 162-169, 1199-1207
- Jeffrey, Jack, see Pejchar, Liba, and —
- Jenkinson AOU Service Award, 2003: Peter Stettenheim, 265-266
- Johnson, Ned Keith, 1932-2003, In Memoriam, 600-602
- Jones, Andrew W., see Zink, Robert M., and —
- Jones, Jason, et al., Minimum estimates of survival and population growth for Cerulean Warblers (*Dendroica cerulea*) breeding in Ontario, Canada, 15-22
- Junco hyemalis*, 162-169, 1269-1281
- J. phaenotus*, 404-414
- Junco, Dark-eyed, 162-169, 1269-1281; Yellow-eyed, 404-414
- Kehoe, F. Patrick, see Traylor, Joshua J., et al.
- Keith, G. Stuart, 1931-2003, In Memoriam, 253-256
- Kelly, Elizabeth G., and Forsman, Eric D., Recent records of hybridization between Barred Owls (*Strix varia*) and Northern Spotted Owls (*S. occidentalis caurina*), 806-810
- Kemp, Alan C., see Delpont, Wayne, et al.
- keratinase, 656-659
- Kershner, Eric L., et al., Breeding-season decisions, reneating, and annual fecundity of female Eastern Meadowlarks (*Sturnella magna*) in southeastern Illinois, 796-805
- Kershner, Eric L., et al., Postfledging movements and survival of juvenile Eastern Meadowlarks (*Sturnella magna*) in Illinois, 1146-1154
- Kershner, Eric L., see Walk, Jeffery W., et al.
- Kestrel, American, 738-751, 837-846, 904-916, 1081-1093, 1187-1198; Eurasian, 103-109
- Ketupa zeylonensis*, 35-45
- Keyser, Amber J., et al., Life-history variation and demography in Western Bluebirds (*Sialia mexicana*) in Oregon, 118-133
- Keyser, Marilynne T., see Keyser, Amber J., et al.
- Keyser, Patrick D., see Hamel, Paul B., et al.
- Kingbird, 1081-1093; Cassin's, 1081-1093; Eastern, 1103-1121; Tropical, 1005-1013; Western, 1081-1093; White-throated, 1005-1013
- Kingfisher, 35-45; Belted, 35-45; Pied, 35-45
- Kinglet, Golden-crowned, 162-169; Ruby-crowned, 74-87, 162-169, 289-301, 738-751, 1040-1055
- Kitaysky, Alexander S., see Thompson, Christopher W., and —
- Kitchens, Wiley M., see Dreitz, Victoria J., et al.
- Kite, Black, 985-995; Snail, 894-903; Swallow-tailed, 404-414; White-tailed, 904-916

- Kiure, Jacob, see Beresford, Pamela, et al.
 kiwis, 308–317
Knipolegus aterrimus, 1005–1013
 Knot, Red, 333–344, 1094–1102
 Koenig, Walter D., and Sherman, Paul W., In Memoriam: Frank Alois Pitelka, 1916–2003, 963–965
 Konarzewski, Marek, see Brzek, Paweł, and —
 Kosciuch, Karl L., and Langerhans, R. Brian, Evolution of coloniality via commodity selection: What about variance?, 257–262
 Krapu, Gary L., et al., Patterns of variation in clutch sizes in a guild of temperate-nesting dabbling ducks, 695–706
 Kratter, Andrew W., see Banks, Richard C., et al.
 Kristin, Anton, see Hoi, Herbert, et al.
 Krohn, William B., see Vickery, Peter D., and —
 Kroodsma, Donald E., 263–264
 Kulikova, Irina V., et al., Asymmetric hybridization and sex-biased gene flow between Eastern Spot-billed Ducks (*Anas zonorhynchos*) and Mallards (*A. platyrhynchos*) in the Russian Far East, 930–949
 Kunc, Hansjoerg P., see Amrhein, Valentin, et al.
 Kwit, Charles, et al., Fruit abundance and local distribution of wintering Hermit Thrushes (*Catharus guttatus*) and Yellow-rumped Warblers (*Dendroica coronata*) in South Carolina, 46–57
 Lafón-Terrazas, Alberto, see Macías-Duarte, Alberto, et al.
Lagonosticta rhodopareia, 156–161
L. senegala, 156–161
Lagopus lagopus, 985–995
L. leucura, 985–995
L. leucurus, 985–995
L. muta, 985–995
L. mutus, 985–995, 1238–1249
L. scotica, 985–995
L. scoticus, 985–995
 LaHaye, William S., et al., Temporal variation in the vital rates of an insular population of Spotted Owls (*Strix occidentalis occidentalis*): Contrasting effects of weather, 1056–1069
Lampornis calolaema, 985–995
L. calolaemus, 985–995
L. castaneiventris, 985–995
Lamprospiza melanoleuca, 88–102
Lamprotornis nitens, 435–445
L. superbus, 435–445
 landbirds, 865–876, 1040–1055
 landscape mosaic, 192–202
 Langerhans, R. Brian, see Kosciuch, Karl L., and —
 Laniidae, 1155–1160
Lanio fulvus, 88–102
Lanius collaris, 1226–1237
L. collurio, 361–371, 1155–1160
L. cristatus, 35–45
L. excubitor, 1155–1160
L. ludovicianus, 435–445, 1081–1093
L. minor, 557–564
 Lank, David B., see Evans Ogden, Lesley J., et al.
 Lanyon, Scott M., see Price, J. Jordan, and —
 Laridae, 380–390
 Lark, Horned, 1081–1093, 1130–1145; Sky, 796–805
Larus, 243–252
L. atlanticus, 243–252
L. californicus, 950–962
L. delawarensis, 170–177, 950–962
L. dominicanus, 243–252
L. fuscus, 243–252
L. glaucescens, 380–390
L. ridibundus, 361–371
L. schistisagus, 452–462
L. scoresbii, 243–252
Larvivora sibilans, 985–995
 Lawes, Michael J., see Pryke, Sarah R., and —
 Laybourne, Roxie Collie, 1910–2003, In Memoriam, 1282–1285
 laying stage, 587–599
 Leedy, Daniel Loney, 1912–2003, In Memoriam, 972
Legatus leucophaeus, 88–102
Leiothrix argentea, 1155–1160
Leipoa ocellata, 308–317
Lepidocolaptes albolineatus, 88–102
L. fuscus, 1155–1160
Leucopternis albigollis, 88–102, 985–995
L. melanops, 88–102
L. plumbea, 985–995
L. plumbeus, 985–995
L. polionota, 985–995
L. polionotus, 985–995
L. schistacea, 985–995
L. schistaceus, 985–995
L. semiplumbea, 985–995
L. semiplumbeus, 985–995
 Levey, Douglas J., see Borgmann, Kathi L., et al.
 Levey, Douglas J., see Jahn, Alex E., et al.
 Levey, Douglas J., see Kwit, Charles, et al.
 life-history, — evolution, 289–301; — variation, 118–133
Limnolophus swainsonii, 788–795
Limosa lapponica, 1094–1102
 Lindell, Catherine A., see Cohen, Emily B., and —
Liocichla phoenicea, 435–445
Lipaugus subalaris, 1155–1160
L. vociferans, 88–102
 lipid reserves, 695–705, 917–929
 Liversidge, Richard, 1926–2003, In Memoriam, 603
 locomotor activity, 1187–1198
Locustella luscinioides, 35–45
 Logan, Clara J., see Henson, Shandelle M., et al.
 logging, 162–169, 1070–1080
 logistic-exposure model, 526–540
Lonchura bicolor, 435–445

- Long, Jennifer A., and Holberton, Rebecca L.,
Corticosterone secretion, energetic condition,
and a test of the migration modulation
hypothesis in the Hermit Thrush (*Catharus
guttatus*), a short-distance migrant, 1094–1102
longevity, 302–307
Longspur, Chestnut-collared, 1081–1093, 1130–1145;
Lapland, 1238–1249; Smith's, 446–451
loons, 35–45
Loos, Elizabeth R., and Rohwer, Frank C., Laying-
stage nest attendance and onset of incubation in
prairie nesting ducks, 587–599
Lophodytes cucullatus, 148–155
Lophornis brachylopha, 985–995
L. brachylophus, 985–995
Lovette, Irby J., Mitochondrial dating and mixed
support for the "2% rule" in birds, 1–6
Lovette, Irby J., review by, 1298–1299
Lovette, Irby J., see Sweeney, Robert J., et al.
Luscinia, 23–34
L. megarhynchos, 110–117, 1155–1160
L. sibilans, 985–995
lyrebirds, 1155–1160

Machetornis rixosa, 985–995
M. rixosus, 985–995, 1155–1160
Macías-Duarte, Alberto, et al., Reproduction, prey,
and habitat of the Aplomado Falcon (*Falco
femorialis*) in desert grasslands of Chihuahua,
Mexico, 1081–1093
Macrocephalon maleo, 308–317
macroecology, 604–609
Macronectes giganteus, 243–252, 422–434
Magpie, Black-billed, 950–962; Eurasian, 361–371,
372–379
Majid, Samia, see Goldstein, Gerald, et al.
Malau, 308–317
Maleo, 308–317
Mallard, 497–508, 526–540, 587–599, 695–706, 917–
929, 930–949
Malleefowl, 308–317
Maluridae, 1155–1160
Malurus leucopterus, 652–655
M. splendens, 1155–1160
Manacus aurantiacus, 985–995
M. manacus, 1155–1160
M. vitellinus, 985–995
M. v. viridiventris, 985–995
management, 7–14, 58–67, 134–147, 192–202,
243–252, 380–390, 515–525, 738–751, 894–903,
1031–1039
manakins, 88–102
Manegold, Albrecht, et al., Miocene songbirds and
the composition of the European passeriform
avifauna, 1155–1160
Marchant, Stephen, 1912–2003, In Memoriam, 971
marine specialization, 752–766
MARK, 707–716, 865–876

Martin, Fairy, 302–307; House, 1187–1198; Sand,
985–995; Southern, 1005–1013
Martin, Jean-Louis, see Tourenq, Christophe, et al.
Martin, Thomas E., Avian life-history evolution has
an eminent past: Does it have a bright future?,
289–301
Martínez-Meyer, Enrique, see Nakazawa, Yoshinori,
et al.
mass-corrected molt score, 372–379
Massemín, Sylvie, see Durant, Joël M., et al.
mate, — advertisement, 486–496; — choice, 156–161
material resources hypothesis, 877–893
maternal origin, 156–161
mating systems, 308–317, 391–403, 446–451, 473–485
Maute, Kimberly L., see Sieving, Kathryn E., et al.
Mayfield logistic regression, 707–716
Mayfield's estimator, 134–147, 526–540, 707–716
Mayr, Gerald, see Manegold, Albrecht, et al.
Mays, Herman L., Jr., see Byers, Bruce E., et al.
Mazar Barnett, Juan, see Herzog, Sebastian K., and —
McCarty, John P., see Kwit, Charles, et al.
McCracken, Kevin G., see Kulikova, Irina V., et al.
McFarland, Kent P., see Strong, Allan M., et al.
Meadowlark, 224–229, 1081–1093; Eastern, 767–776,
796–805, 1081–1093, 1146–1154; Western, 134–
147, 824–836, 1081–1093, 1130–1145
Megaceryle, 985–995
Megadyptes, 422–434
M. antipodes, 422–434
Megapode, Melanesian, 308–317; New Guinea, 308–
317; Nicobar, 308–317; Orange-footed, 308–317;
Polynesian, 302–307, 308–317
Megapodius decollatus, 308–317
M. nicobariensis, 308–317
M. pritchardii, 302–307, 308–317
M. reinwardt, 308–317
Megarhynchus pitangua, 1155–1160
Megascops asio, 738–751
Melanerpes formicivorus, 1081–1093
M. pygmaeus, 985–995
M. radiolatus, 985–995
M. rubricapillus, 985–995
melanin, 652–655, 656–659
Melanitta, 950–962
M. fusca deglandi, 950–962
Meleagris gallopavo, 148–155, 717–725
Meliphagidae, 1155–1160
Melospiza lincolni, 162–169
M. melodia, 391–403, 404–414, 435–445, 652–655, 738–
751, 824–836, 877–893
M. m. fallax, 656–659
M. m. morphna, 656–659
Melozone *biarcuata*, 985–995
M. biarcuatum, 985–995
M. kieneri, 985–995

- Melvin, Edwin F., and Parrish, Julia, Seabird Bycatch: Trends, Roadblocks, and Solutions (rev.), 627-628
- Menura novaehollandiae*, 1155-1160
- Menuridae, 1155-1160
- Merganser, Hooded, 148-155
- Merkel, Friedrich Wilhelm "Fritz", 1911-2002, in Memoriam, 1291
- Merlin, 1081-1093
- Meropidae, 811-823, 1187-1198
- Merops apiaster*, 541-547, 811-823
- M. bullockoides*, 811-823
- M. bullocki*, 811-823
- M. nubicoides*, 811-823
- M. orientalis*, 811-823
- M. ornatus*, 811-823
- M. persicus*, 811-823
- M. pusillus*, 811-823
- M. viridis*, 811-823
- Mesléard, François, see Tourenq, Christophe, et al.
- Mesophoyx intermedia*, 985-995
- meta-analysis, 1056-1069
- Metopidius*, 391-403
- M. indicus*, 391-403
- Mezey, Jason G., see James, Frances C., and —
- Micrastur gilvicollis*, 88-102
- microbial ectoparasites, 1262-1268
- Microparra capensis*, 391-403
- microsatellite DNA, 788-795
- migrant, short-distance, 46-57, 1094-1102
- migration modulation response, 1094-1102
- migration, 811-823, 837-846, 1094-1102; austral, 1005-1013; New World, 1005-1013; partial, 1005-1013; spring, 917-929
- migratory, — fattening, 1094-1102; — movements, 837-846; — systems, 610-618
- Miliaria calandra*, 1226-1237
- Milvago chimachima*, 88-102
- Milvus migrans*, 985-995
- Mimus polyglottos*, 1081-1093, 1155-1160
- M. triurus*, 1005-1013
- Mino anais*, 1155-1160
- Miocene songbirds, 1155-1160
- Miocorvus larteti*, 1155-1160
- Mionectes macconnelli*, 88-102
- Mississippi Flyway, 917-929
- mitochondrial, — dating, 1-6; — DNA, 1-6, 23-34, 156-161, 224-229, 230-242, 318-332, 660-681, 682-694, 847-864, 930-949, 1014-1018, 1019-1030; — genes, 752-766, 1031-1039
- Mitu, 682-694
- M. mitu*, 682-694
- M. salvini*, 682-694
- M. tomentosa*, 682-694
- M. tuberosa*, 682-694
- mixed-grass prairie, 1130-1145
- Miyaki, Cristina Y., see Tavares, Erika S., et al.
- Mlíkovsky, Jiri, Cenozoic Birds of the World, Part 1: Europe (rev.), 623-627
- Mniotilta varia*, 738-751, 1040-1055
- mobbing, 738-751
- Mock, Douglas W., 264-265
- Mockingbird, Northern, 1081-1093; White-banded, 1005-1013
- molecular, — clock, 1-6, 318-332; — phylogeny, 847-864; — systematics, 318-332
- mollusk shells, 148-155
- Molothrus aeneus*, 1172-1186
- M. ater*, 7-14, 308-317, 354-360, 726-737, 767-776, 824-836, 865-876, 877-893, 1081-1093, 1103-1121, 1130-1145, 1172-1186
- M. bonariensis*, 1172-1186
- M. oryzivora*, 1172-1186
- molt, 372-379; flight-feather, 23-34; primary, 23-34; secondary, 23-34; — score index, 372-379
- Monaghan, Pat, see Hilton, Geoff M., et al.
- Monasa atra*, 88-102
- Monjita, Black-crowned, 1005-1013
- monogamous male, 1226-1237
- monogamy, genetic, 302-307, 308-317; social, 302-307, 308-317, 788-795, 811-823
- montane, — avifauna, 23-34; — forest, 23-34; — breeding environment, 446-451
- montane-breeding birds, 1269-1281
- Montevicchi, W. A., review by, 627-628
- Monticola solitarius*, 985-995
- M. s. philippensis*, 985-995
- Montoya, Angel B., see Macias-Duarte, Alberto, et al.
- Moore, Jeffrey E., et al., Field comparison of removal and modified double-observer modeling for estimating detectability and abundance of birds, 865-876
- moorhens, 1122-1129
- Morphnus guianensis*, 88-102
- morphological features, 973-976
- morphology, 422-434, 565-586, 847-864, 1031-1039, 1155-1160; external, 23-34
- morphometric, — characters, 660-681; — traits, 178-183
- Morrison, R. I. Guy, and Hobson, Keith A., Use of body stores in shorebirds after arrival on High-Arctic breeding grounds, 333-344
- Morrison, Scott A., et al., Annual survivorship of the sedentary Rufous-crowned Sparrow (*Aimophila ruficeps*): No detectable effects of edge or rainfall in southern California, 904-916
- mortality, 526-540; age-specific, 289-301; — patterns, 1146-1154
- Morton, Martin L., see Hahn, Thomas P., et al.
- mosaic hybridization, 1014-1018; — model, 565-586
- Motacilla*, 435-445, 682-694
- M. alba*, 1155-1160
- M. angarensis*, 985-995
- M. atricapilla*, 985-995
- M. citreola*, 777-787
- M. curruca*, 985-995
- M. flava*, 435-445, 985-995

- M. macronyx*, 985–995
M. plexa, 985–995
M. simillima, 985–995
M. striata, 985–995
M. taivana, 985–995
M. tschutschensis, 985–995
M. zaisanensis, 985–995
 Mourer-Chauviré, Cécile, review by, 623–627
 Mourer-Chauviré, Cécile, see Manegold, Albrecht, et al.
 movement patterns, 162–169
 Mukhin, Andrey, Night movements of young Reed Warblers (*Acrocephalus scirpaceus*) in summer: Is it postfledging dispersal?, 203–209
 Mumme, Ronald L., see Kwit, Charles, et al.
 Murre, 148–155; Common, 35–45, 148–155, 1161–1171; Thick-billed, 148–155, 452–462
 Murrelet, Kittlitz's, 35–45; Marbled, 1161–1171; Xantus's, 1161–1171
Muscicapa dauurica, 985–995
M. griseisticta, 985–995
M. sibirica, 985–995
M. striata, 985–995
 Muscicapidae, 88–102, 1155–1160
 Muscicapinae, 23–34
Muscisaxicola, 752–766
M. capistrata, 1005–1013
Myiarchus cinerascens, 1081–1093
M. crinitus, 738–751, 1040–1055
M. ferox, 88–102
M. swainsoni, 610–618, 1005–1013
M. tuberculifer, 88–102
M. tyrannulus, 435–445
Myioborus, 985–995
Myiodynastes maculatus, 88–102
Myiopagis caniceps, 88–102
M. gaimardii, 88–102
Myornis, 1155–1160
Myrmeciza longipes, 486–496
Myrmotherula, 88–102
M. brachyura, 88–102, 985–995
M. h. ignota, 985–995
M. fulviventris, 486–496
M. ignota, 985–995
M. obscura, 985–995
Myzomela rubrata, 1155–1160
 Naguib, Marc, see Amrhein, Valentin, et al.
 Naka, Luciano Nicolás, Structure and organization of canopy bird assemblages in central Amazonia, 88–102
 Nakazawa, Yoshinori, et al., Seasonal niches of Nearctic-Neotropical migratory birds: Implications for the evolution of migration, 610–618
 native-prairie patches, 1130–1145
 Navarro-Sigüenza, Adolfo G., see Nakazawa, Yoshinori, et al.
 Nearctic-Neotropical species, 610–618
Nectarinia, 23–34
N. afra, 660–681
N. asiatica, 1155–1160
N. chalybea, 660–681
N. chloropygia, 660–681
N. fuelleborni, 660–681
N. loveridgei, 660–681
N. mediocris, 660–681
N. m. bensoni, 660–681
N. m. fuelleborni, 23–34, 660–681
N. m. mediocris, 660–681
N. m. usambarica, 660–681
N. minulla, 660–681
N. moreaui, 23–34, 660–681
N. preussi, 660–681
N. regia, 660–681
N. rufipennis, 660–681
N. sperata, 435–445
N. usambarica, 660–681
 Nectariniidae, 660–681, 1155–1160
 nectarivores, 88–102
 Neoaves, 230–242
Neolestes torquatus, 1155–1160
 Neotropical, — avifauna, 318–332, 1005–1013; — migrant, 1019–1030; — resident, 404–414
 nest, — attendance, 587–599; — characteristics, 1226–1237; — concealment, 361–371, 950–962; — construction, 777–787; — dispersion, 1226–1237; — failure, 134–147, 1250–1261; — microclimate, 777–787; — placement, 767–776; — predation, 162–169, 289–301, 361–371, 446–451, 877–893, 1214–1225, 1226–1237; — predators, 162–169; — sites, 1250–1261; — success, 134–147, 354–360, 361–371, 497–508, 526–540, 707–716, 726–737, 1070–1080, 1122–1129, 1214–1225; — survival models, 134–147; — survival, 526–540, 767–776, 950–962
 nesting, — ecology, 148–155, 354–360, 950–962; — habitat, 243–252, 404–414; — material, 777–787; — period, 58–67
 nest-site, — characteristics, 796–805, 1250–1261; — effects, 726–737; — selection, 950–962; — suitability, 1081–1093
 nest-survival analysis, 707–716
Netta rufina, 637–651
 Neudorf, Diane L. H., Extrapair paternity in birds: Understanding variation among species, 302–307
 new species, 1031–1039
 Newton, Ian, see Dawson, Alistair, and —
 Newton, Ian, The Speciation and Biogeography of Birds (rev.), 1296–1298
 niche specialization, 548–556
 niches, ecological, 610–618; seasonal, 610–618
 Niemi, Gerald J., review by, 980–982
 night movements, 203–209
 Nighthawk, 1081–1093; Common, 1081–1093; Lesser, 1081–1093

- Night-Heron, Black-crowned, 192–202
 Nightingale, 110–117
 Nightjar, Gray, 985–995; Jungle, 985–995
 Nixon, Wendy A., see Sinclair, Pamela H., et al.
 nocturnal activity, 203–209
 nomenclature, 985–995
 nonbreeding season, 486–496
 nonheritable variation, 345–353
 nonmigratory habit, 904–916
 nonpasseriform, 1155–1160
 nonpasserines, 88–102
 north-temperate systems, 289–301
Notharchus macrorhynchus, 88–102
N. tectus, 88–102
Nothocrax, 682–694
N. urumutum, 682–694
 nuclear DNA, 23–34
 Nudds, Thomas D., see Drever, Mark C., et al.
Numida galeata, 985–995
N. galeatus, 985–995
N. meleagris, 985–995
N. mitrata, 985–995
N. mitratus, 985–995
 nutrient limitation hypothesis, 587–599; — reserves, 917–929
 nutrients, endogenous, 333–344; exogenous, 333–344
 nutritional, — requirements, 463–472; — resources, 333–344
 Nyctibiidae, 88–102
Nyctibius grandis, 88–102
N. leucopterus, 88–102
Nycticorax nycticorax, 192–202
- O'Connor, Raymond J., and Wagner, Tansy L.,
 A test of a regression-tree model of species
 distribution, 604–609
 O'Loughlin, Adrian L., and Rothstein, Stephen
 I., Divergent sexual responses to different
 categories of foreign courtship songs in female
 Brown-headed Cowbirds (*Molothrus ater*),
 824–836
- Oceanites oceanicus*, 422–434
 Oceanitinae, 1–6
Oceanodroma leucorhoa, 422–434
Ocyalus, 224–229
O. latirostris, 224–229
Oenanthe, 23–34
 oligopeptides, 656–659
 omnivores, 88–102
 onset of egg laying, 1269–1281
Oporornis agilis, 1040–1055
O. formosus, 726–737, 1040–1055
O. philadelphia, 1040–1055
 Oren, David C., see Whitney, Bret M., et al.
Oreomystis bairdii, 1–6
O. mana, 548–556
Oreopelia leucometopius, 985–995
 Oriole, Baltimore, 1040–1055; Scott's, 1081–1093
 orioles, 224–229
Oriolus auratus, 435–445
O. oriolus, 1155–1160
Ornithion inermis, 88–102
 ornithology, 637–651, 973–976
 Oropendola, Green, 224–229; Montezuma, 224–229;
 Olive, 224–229
Ortalis canicollis, 682–694
 Orthonychidae, 1155–1160
Orthonyx spaldingi, 1155–1160
Orthopsittacus, 230–242
 oscines, 1155–1160
 osteology, 1155–1160
 Otter, Ken A., see Fort, Kevin T., and —
Otus asio, 46–57
O. scops, 541–547
 ovary analysis, 103–109
 Ovenbird, 726–737, 738–751, 752–766, 1040–1055
 Overview, 7–14, 302–307, 652–655, 1014–1018
 overwintering, 74–87
 ovulation, 103–109
- Owl, 35–45; Barn, 103–109; Barred, 806–810; Brown
 Fish, 35–45; California Spotted, 1056–1069;
 Eagle, 35–45; Eurasian Scops, 541–547; Great
 Horned, 950–962, 1081–1093; Mexican Spotted,
 1056–1069; Northern Spotted, 806–810, 1056–
 1069; Rufous-legged, 515–525; Snowy, 35–45;
 Spotted, 1056–1069; Tawny, 541–547, 1056–1069;
 Tengmalm's, 1103–1121
- Pachycephala lanioides*, 1155–1160
P. rufiventris, 1155–1160
Pachyptila desolata, 422–434
Pachyramphus marginatus, 88–102
P. minor, 88–102
P. surinamus, 88–102
Padda oryzivora, 435–445
 pair bond, 308–317
 Palmer, Ralph Simon, 1914–2003, In Memoriam,
 966–968
- Panurus biarmicus*, 302–307
Panyptila cayannensis, 88–102
Paradisaea rubra, 1155–1160
 Paradisaidae, 1155–1160
 paradise whydahs, 156–161
 Pardalotidae, 1155–1160
Pardalotus substriatus, 1155–1160
 parental, — care, 302–307, 308–317, 1199–1207;
 — investment, 391–403, 1238–1249;
 — provisioning, 452–462, 463–472
 Paridae, 435–445, 738–751, 1155–1160
Paroaria coronata, 435–445
Paroreomyza montana, 1–6
Parotia lawesii, 391–403
 Parotia, Lawes', 391–403
 Parr, Michael J., see Chipley, Robert M., et al.
 Parrish, Julia K., see Hamel, Nathalie J., et al.
 Parrish, Julia, see Melvin, Edwin F., and —

- Parrot, 35–45, 88–102, 230–242, 318–332, 1214–1225;
Lilac-crowned, 1214–1225; Red-crowned, 1214–1225
- Parrotlet, Green-rumped, 717–725
- partridges, 1–6
- Parula americana*, 435–445, 738–751, 1040–1055
- Parula*, Northern, 738–751, 1040–1055
- Parulidae, 788–795, 1005–1013
- parulids, 15–22
- Parulinae, 88–102
- Parus*, 289–301
- P. caeruleus*, 435–445, 652–655, 1155–1160
- P. major*, 184–191, 354–360, 1155–1160
- P. palustris*, 404–414
- Pashley, David N., see Chipley, Robert M., et al.
- Pasquet, Eric, see Austin, Jeremy J., et al.
- Passer, 435–445
- P. domesticus*, 118–133, 302–307, 435–445, 837–846, 1155–1160, 1187–1198, 1226–1237
- P. griseus*, 435–445
- Passerculus sandwichensis*, 473–485, 1081–1093, 1130–1145
- Passerella iliaca*, 162–169, 985–995, 1269–1281
- Passeridae, 435–445, 1155–1160
- Passeriformes, 354–360, 752–766, 1155–1160
- Passerina amoena*, 877–893
- P. caerulea*, 1081–1093
- P. cyanea*, 435–445, 726–737, 865–876, 1040–1055, 1103–1121
- passerines, 88–102, 118–133, 372–379, 435–445, 637–651, 738–751, 752–766, 777–787, 877–893, 1103–1121, 1130–1145, 1146–1154, 1155–1160, 1199–1207
- patch, — shape, 1130–1145; — size, 1130–1145
- Pauxi*, 682–694
- P. pauxi*, 682–694
- P. unicornis*, 682–694
- Payne, Robert B., and Sorenson, Michael, D., Behavioral and genetic identification of a hybrid *Vidua*: Maternal origin and mate choice in a brood-parasitic finch, 156–161
- Peak, Rebecca G., et al., Factors affecting songbird nest survival in riparian forests in a midwestern agricultural landscape, 726–737
- Pearson, Scott F., see Borgmann, Kathi L., et al.
- Pearson, Scott F., see Kwit, Charles, et al.
- Pechacek, Peter, Spacing behavior of Eurasian Three-toed Woodpeckers (*Picoides tridactylus*) during the breeding season in Germany, 58–67
- perched song, 824–836
- Peer, Brian D., and Sealy, Spencer G., Fate of grackle (*Quiscalus* spp.) defenses in the absence of brood parasitism: Implications for long-term parasite-host coevolution, 1172–1186
- Pejchar, Liba, and Jeffrey, Jack, Sap-feeding behavior and tree selection in the endangered Akiapolaau (*Hemignathus munroi*) in Hawaii, 548–556
- Pelecanoides urinatrix*, 422–434
- Pelecanoididae, 35–45
- Pelecanus erythrorhynchos*, 950–962
- Pelican, American White, 950–962
- Penelope marail*, 88–102
- P. obscura*, 682–694
- Penguin, 148–155, 422–434; Adelie, 148–155; King, 1161–1171; Macaroni, 148–155; Magellanic, 148–155, 243–252
- Pereira, Sérgio Luiz, and Baker, Allan J., Vicariant speciation of curassows (Aves, Cracidae): A hypothesis based on mitochondrial DNA phylogeny, 682–694
- Perisoreus canadensis*, 162–169, 435–445
- P. infaustus*, 162–169, 1199–1207
- Perkins, Kelly A., et al., Flushing, capture, and bleeding do not affect return rate of female Wood Thrushes (*Hylocichla mustelina*) in Delaware, 354–360
- Perspectives in Ornithology, 1–6, 289–301, 637–651, 1005–1013
- Peterson, A. Townsend, see Nakazawa, Yoshinori, et al.
- Petrel, 35–45, 422–434, 847–864; Dark-rumped, 985–995; Galapagos, 985–995; Great-winged, 985–995; Southern Giant, 243–252
- Petrochelidon fulva*, 1122–1129
- P. pyrrhonota*, 1122–1129
- Petroica traversi*, 354–360
- Petroicidae, 1155–1160
- Pewee, Crescent-eyed, 985–995
- Phaethornis*, 88–102
- Phalacrocoracidae, 148–155
- Phalacrocorax aristotelis*, 103–109
- P. atriceps*, 243–252
- P. brasiliensis*, 243–252
- P. magellanicus*, 243–252
- P. penicillatus*, 985–995
- Phalaenoptilus nuttallii*, 1081–1093
- Phalarope, Grey, 391–403; Red-necked, 391–403
- Phalaropus fulicarius*, 391–403
- P. lobatus*, 391–403
- Pharomachrus pavoninus*, 88–102
- Phasianidae, 515–525
- Phasianus colchicus*, 308–317
- Pheasant, 515–525; Ring-necked, 308–317
- Pheucticus ludovicianus*, 435–445, 1040–1055
- P. melanocephalus*, 1081–1093
- Philepitta castanea*, 1155–1160
- Philepittidae, 1155–1160
- Philetairus socius*, 1199–1207
- Phillips, Richard A., see Catry, Paulo, et al.
- philopatry, natal, 1103–1121
- philosophy of biology, 637–651
- Philydor erythrocerum*, 985–995
- P. erythrocerus*, 985–995
- P. erythronotus*, 985–995
- P. erythronotus*, 985–995
- P. fuscipennis*, 985–995

- P. fuscipennis*, 985–995
P. rufum, 985–995
P. rufus, 985–995
 phloem sap, 548–556
Phoebastria, 1–6
P. irrorata, 847–864
Phoebe, Eastern, 738–751; Say's, 1081–1093
Phoebetria palpebrata, 422–434
Phoenicopterus ruber, 541–547
 Phoeniculidae, 178–183
Phoeniculus purpureus, 178–183
Phylloscartes, 88–102
P. virescens, 88–102
Phylloscopus, 877–893
P. borealis, 985–995
P. collybita, 110–117
P. sibilatrix, 985–995
P. trochilus, 302–307, 985–995, 1238–1249, 1155–1160
P. t. acredula, 985–995
P. t. trochilus, 985–995
 phylogenetic, — analysis, 318–332, 660–681;
 — distribution, 435–445; — hypothesis, 435–445;
 — relationships, 224–229, 230–242, 318–332,
 435–445, 682–694, 752–766, 847–864, 930–949,
 1155–1160
 phylogeny, 422–434
 physiological condition, 203–209
Phytotoma rara, 1155–1160
Piaya melanogaster, 88–102
Pica, 985–995
P. hudsonia, 950–962
P. pica, 361–371, 372–379, 985–995, 1155–1160
Picathartes oreas, 1155–1160
 Picarthidae, 1155–1160
 Picidae, 88–102
Picoides, 985–995
P. arcticus, 58–67, 509–514
P. borealis, 58–67, 74–87, 1103–1121
P. dorsalis, 58–67
P. major, 58–67, 509–514
P. medius, 58–67, 509–514
P. minor, 509–514
P. pubescens, 74–87, 509–514, 738–751
P. scalaris, 1081–1093
P. tridactylus, 58–67
P. villosus, 509–514
P. viridis, 509–514
Piculus chrysocloros, 88–102
P. flavigula, 88–102
Picumnus exilis, 88–102
 Pigeon, 88–102; Chilean, 515–525; Rock, 103–109
Pinguinus impennis, 35–45
 Pintail, Northern, 497–508, 587–599, 695–706
Pionopsitta caica, 88–102
Pionus, 230–242
P. fuscus, 88–102
P. menstruus, 88–102, 230–242
Pipilo chlorurus, 289–301, 404–414, 1081–1093, 1269–1281
P. erythrophthalmus, 74–87, 738–751, 824–836
P. fuscus, 435–445
P. maculatus, 1014–1018, 1081–1093
P. ocai, 1014–1018
 Pipit, American, 1081–1093, 1250–1261; Richard's,
 35–45; Sprague's, 1081–1093, 1130–1145; Water,
 777–787
Pipra erythrocephala, 88–102, 1155–1160
P. mentalis, 435–445
P. pipra, 88–102
 Pipridae, 88–102
Piprites chloris, 88–102
Piranga olivacea, 435–445, 1040–1055
Pitangus sulphuratus, 1155–1160
 Pitelka, Frank Alois, 1916–2003, In Memoriam,
 963–965
 Pitohui, 1262–1268
 Pitta, 435–445
P. granatina, 1155–1160
P. guajana, 1155–1160
P. moluccensis, 435–445
P. sordida, 1155–1160
 Pitta, 1155–1160; Garnet, 1155–1160
 Pittidae, 1155–1160
Pitylus grossus, 88–102
 plant biomass, 515–525
 Pleistocene refugia theory, 1–6
 Ploceidae, 435–445
 Ploceinae, 1226–1237
Ploceus, 1172–1186
P. castanops, 777–787
P. cucullatus, 1172–1186
P. ocularis, 435–445
P. velatus, 435–445
 Plover, Mongolian, 985–995; Mountain, 526–540
 plumage, — characters, 660–681; — phenotype,
 1019–1030
 Pochard, Red-crested, 637–651
Podargus ocellatus, 541–547
 Podicipedidae, 35–45, 777–787
Poecile atricapillus, 435–445, 1040–1055, 1070–1080
P. carolinensis, 74–87, 738–751, 865–876
P. gambeli, 777–787
 point-count data, 865–876
Poliophtila caerulea, 435–445, 738–751, 1040–1055
P. guianensis, 88–102
P. melanura, 1081–1093
 Polioptilinae, 88–102, 435–445
 polyandry, 308–317, 391–403
Polyboroides typus, 1226–1237
 Polyerata, 985–995
 polygynandry, 446–451
 polygynous male, 1226–1237
 polygyny, 308–317
 Pomatostomidae, 1155–1160
Pomatostomus superciliosus, 1155–1160
 pond density, 497–508
Poocetes gramineus, 1081–1093

- Poorwill, Common, 1081–1093
 population, — biology, 796–805; — change, 1056–1069; — decline, 15–22, 917–929, 950–962, 1019–1030; — dynamics, 1146–1154; — growth, 118–133; — trends, 7–14
- Porzana carolina*, 1081–1093
 post-egg period, 148–155
 postfledging, — movements, 203–209, 1146–1154; — period, 1146–1154
- Pough, Richard H., 1904–2003, In Memoriam, 969
- prairie, — environments, 497–508; — pothole landscape, 695–705
- precipitation, 695–705, 1081–1093
- precocial species, 308–317
- predation, 806–810; — hypothesis, 877–893; — risk, 134–147, 738–751
- predator management, 497–508
- pre migratory period, 203–209
- Prescott Bluebird Recovery Project, 118–133
- prey, — abundance, 894–903, 1081–1093; — biomass, 446–451
- Price, J. Jordan, and Lanyon, Scott M., Song and molecular data identify congruent but novel affinities of the Green Oropendola (*Psarocolius viridis*), 224–229
- Primack, Richard B., Essentials of Conservation Biology, 3rd ed. (rev.), 980–982
- Primolius*, 230–242
P. auricollis, 230–242
P. coultoni, 230–242
P. maracana, 230–242
- prions, 422–434
- Procellaria aequinoctialis*, 422–434
P. macroptera, 985–995
P. m. gouldi, 985–995
- Procellariidae, 1–6
 Procellariiformes, 1–6, 35–45, 422–434, 847–864
Procnias nudicollis, 1155–1160
P. tricarunculata, 985–995
P. tricarunculatus, 985–995
- productivity, 1081–1093
- Progne*, 88–102
P. chalybea, 88–102
P. modesta, 1005–1013
P. subis, 88–102
- Promislow, Daniel E. L., see Keyser, Amber J., et al.
- Protonaria citrea*, 354–360, 1103–1121
- provisioning, 1161–1171
- Prunella*, 1155–1160
P. collaris, 184–191
P. modularis, 446–451
- Pryke, Sarah R., and Lawes, Michael J., Female nest dispersion and breeding biology of polygynous Red-collared Widowbirds (*Euplectes ardens*), 1226–1237
- Psaltiriparus minimus*, 435–445
Psarisomus dalhousiae, 1155–1160
Psarocolius, 224–229
P. angustifrons alfredi, 224–229
P. a. angustifrons, 224–229
P. a. atrocastaneus, 224–229
P. a. oleagineus, 224–229
P. atrovirens, 224–229
P. decumanus, 224–229
P. d. decumanus, 224–229
P. d. insularis, 224–229
P. d. maculosus, 224–229
P. d. melanterus, 224–229
P. oseryi, 224–229
P. viridis, 88–102, 224–229
P. wagleri, 224–229
P. w. ridgwayi, 224–229
P. w. wagleri, 224–229
- Pseudocolaptes boissonneautii*, 1155–1160
- Psittacidae, 88–102, 230–242, 1214–1225
- Psittaciformes, 35–45, 230–242, 717–725
- Psittacinae, 230–242, 1214–1225
- Ptarmigan, Rock, 1238–1249
- Pterodroma*, 847–864
P. cahow, 847–864
P. externa, 847–864
P. feae-madeira, 847–864
P. hasitata, 847–864
P. incerta, 422–434, 847–864
P. inexpectata, 847–864
P. lessoni, 847–864
P. macroptera, 847–864, 985–995
P. neglecta, 847–864, 985–995
P. phaeopygia, 847–864, 985–995
- Pteroglossus, 230–242
P. viridis, 88–102
- Pterotochos megapodius*, 1155–1160
- P. tarnii*, 515–525
- Ptilonorhynchidae, 1155–1160
- Ptilonorhynchus violaceus*, 1155–1160
- Ptychoramphus*, 985–995
P. aleuticus, 452–462
- puffbirds, 88–102
- Puffin, Atlantic, 35–45, 452–462, 463–472; Horned, 35–45; Tufted, 35–45, 463–472
- Puffinus*, 847–864
P. assimilis, 847–864
P. assimilis-lherminieri, 847–864
P. atrodorsalis, 847–864
P. bailloni, 847–864
P. bannermani, 847–864
P. baroli, 847–864
P. boydi, 847–864
P. bulleri, 847–864
P. carneipes, 847–864
P. colstoni, 847–864
P. creatopus, 847–864
P. dichrous, 847–864
P. elegans, 847–864
P. gavia, 847–864
P. gravis, 422–434, 847–864

- P. griseus*, 422-434, 847-864
P. gunax, 847-864
P. haurakiensis, 847-864
P. heinrothi, 847-864
P. huttoni, 847-864
P. kermadecensis, 847-864
P. lherminieri, 847-864
P. loyemilleri, 847-864
P. mauretanicus, 847-864
P. myrtae, 847-864
P. nativitatis, 847-864
P. newelli, 847-864
P. nicolae, 847-864
P. opisthomelas, 847-864
P. pacificus, 847-864
P. persicus, 847-864
P. polynesiae, 847-864
P. puffinus, 847-864
P. subalaris, 847-864
P. temptator, 847-864
P. tenuirostris, 847-864
P. tunneyi, 847-864
P. yelkouan, 847-864
 Puglisi, Luca, and Adamo, Claudia, Discrimination of individual voices in male Great Bitterns (*Botaurus stellaris*) in Italy, 541-547
 Pycnonotidae, 1155-1160
Pycnonotus jocosus, 1155-1160
 Pygmy-Owl, Austral, 515-525; Eurasian, 541-547
Pygoscelis, 148-155, 422-434
P. adeliae, 422-434
P. antarctica, 422-434
P. papua, 422-434
Pyrocephalus rubinus, 1005-1013, 1081-1093
Pyrrhula pyrrhula, 372-379
Pyrrhuloxia, 1081-1093
Pyrrhura picta, 230-242
Pytilia melba, 156-161
 Quail, Japanese, 170-177, 333-344; Scaled, 1081-1093
 Quail-Dove, Gray-fronted, 985-995; Gray-headed, 985-995; Moustached, 985-995; White-fronted, 985-995
Quelea quelea, 1155-1160
 quiet call, 184-191
Quiscalus, 1172-1186
Q. lugubris, 1172-1186
Q. major, 1172-1186
Q. mexicanus, 1081-1093, 1172-1186
Q. niger, 1155-1160, 1172-1186
Q. quiscula, 435-445, 1172-1186
 Radford, Andrew N., and du Plessis, Morné A., Extreme sexual dimorphism in Green Woodhoopoe (*Phoeniculus purpureus*) bill length: A case of sexual selection?, 178-183
 radiotelemetry, 404-414, 486-496, 796-805, 1146-1154, 1161-1171, 1250-1261, 1269-1281
 radiotracking, 58-67
 radiotransmitters, 1161-1171
 Rail, Virginia, 1081-1093
 rainfall, 904-916
Rallus limicola, 1081-1093
 Ramphastidae, 88-102
Ramphastos tucanus, 88-102
R. vitellinus, 88-102
Ramphocelus carbo, 88-102
 range expansion, 1019-1030
 raptors, 88-102, 103-109, 1056-1069
 Rasmussen, Pamela C., review by, 619-622
 Rasmussen, Pamela C., see Banks, Richard C., et al.
 Ratcliffe, Laurene M., see Tarof, Scott A., and —
 ratites, 1-6
 Raven, Chihuahuan, 1081-1093; Common, 162-169, 361-371, 637-651
 Razorbill, 452-462
 Rebstock, Ginger A., see Boersma, P. Dee, et al.
 recruitment, 68-73, 950-962
 Redpoll, Common, 1238-1249
 Redshank, Common, 372-379, 877-893
 Redstart, 985-995; American, 15-22, 788-795, 877-893, 1040-1055
 refeeding, 1187-1198
 regional variation, 660-681
 regression tree analysis, 604-609
 Regulidae, 1155-1160
Regulus calendula, 74-87, 162-169, 289-301, 435-445, 738-751, 1040-1055
R. regulus, 877-893, 1155-1160
R. satrapa, 162-169
 Reid, Jane M., see Hilton, Geoff M., et al.
 Reid, Sharon, et al., Importance of native bamboo for understory birds in Chilean temperate forests, 515-525
Reinarda squamata, 88-102
Remiz pendulinus, 1155-1160
 removal modeling, 865-876
 Remsen, J. V., Jr., see Banks, Richard C., et al.
 renesting, 796-805, 1250-1261; — rate, 361-371
 Renner, Martin, see Davis, Lloyd S., and —
 Renner, Randy W., see Krapu, Gary L., et al.
 Renton, Katherine, and Salinas-Melgoza, Alejandro, Climatic variability, nest predation, and reproductive output of Lilac-crowned Parrots (*Amazilia finschi*) in tropical dry forest of western Mexico, 1214-1225
 replacement nests, 1250-1261
 reproduction, 1056-1069, 1081-1093
 reproductive, — output, 1214-1225; — potential, 103-109; — rates, 391-403; — role division, 178-183; — state, 110-117; — success, 118-133, 178-183, 257-262, 276-277, 354-360, 446-451, 557-564, 717-725, 788-795, 811-823, 894-903, 917-929, 1122-1129, 1161-1171, 1214-1225, 1238-1249
 residents, 1103-1121

- resource, — abundance, 74–87; — availability, 46–57;
— distribution, 446–451
- retention, 1172–1186
- return rate, 354–360
- Reviewers for *The Auk*, 2004, 1304–1305
- Revis, Hannah C., and Waller, Deborah A.,
Bactericidal and fungicidal activity of ant
chemicals on feather parasites: An evaluation of
anting behavior as a method of self-medication
in songbirds, 1262–1268
- Reynolds, Ronald E., see Krapu, Gary L., et al.
- Rhamphastidae, 230–242
- rhinocryptid, 1155–1160
- Rhinocryptidae, 515–525, 1155–1160
- Rhytipterna simplex*, 88–102
- ribosomal DNA, 230–242
- rice fields, 192–202
- Ridgwayia*, 435–445
- R. pinicola*, 435–445
- Rimmer, Christopher C., see Strong, Allan M., et al.
- Riparia riparia*, 985–995, 1187–1198
- riparian forest, 726–737
- Rising, James D., Geographic Variation in Size and
Shape of Savannah Sparrows (*Passerculus
sandwichensis*) (rev.), 269–270
- Rising, James D., see Banks, Richard C., et al.
- Roadrunner, Greater, 1081–1093
- Robbins, Chandler S., review by, 273–275
- Robertson, Raleigh J., see Jones, Jason, et al.
- Robin, American, 162–169, 354–360, 404–414,
738–751, 865–876, 1103–1121; Chatam Island
Black, 354–360; Rufous-tailed, 985–995; White-
throated, 404–414
- Robson, Craig, Birds of Thailand (rev.), 1299–1300
- Rodewald, Amanda D., review by, 978–980
- Rodewald, Paul G., and Brittingham, Margaret C.,
Stopover habitats of landbirds during fall: Use
of edge-dominated and early-successional
forests, 1040–1055
- Rohwer, Frank C., see Loos, Elizabeth R., and —
Rook, 35–45
- Rosefinch, Scarlet, 361–371
- Rostrhamus sociabilis*, 894–903
- Roth, Roland R., see Perkins, Kelly A., et al.
- Rothstein, Stephen I., see O'Loghlen, Adrian L., and —
Rupicola peruviana, 1155–1160
- R. rupicola*, 391–403
- Ruxton, Graeme D., see Hilton, Geoff M., et al.
- Sadoul, Nicolas, see Tourenq, Christophe, et al.
- Salinas-Melgoza, Alejandro, see Renton, Katherine,
and —
- Sanderling, 333–344
- Sandoz, Alain, see Tourenq, Christophe, et al.
- Sandpiper, Pectoral, 777–787; Purple, 752–766;
Semipalmated, 333–344; Spoonbill, 985–995;
Spoon-billed, 985–995; Spotted, 391–403; White-
rumped, 333–344, 777–787
- Sand-Plover, Greater, 985–995; Lesser, 985–995
- sap feeding, 548–556
- Sapsucker, Yellow-bellied, 509–514, 548–556
- Sarcoramphus papa*, 88–102
- Sargeant, Glen A., see Krapu, Gary L., et al.
- Sargent, Sarah, see Kwit, Charles, et al.
- Saxicola*, 23–34
- S. maura*, 985–995
- S. maurus*, 985–995
- S. rubetra*, 985–995
- S. torquata*, 796–805, 985–995
- S. torquatus*, 985–995
- Saxicolinae, 23–34
- Sayornis phoebe*, 435–445, 738–751
- S. saya*, 1081–1093
- Scaup, Greater, 917–929; Lesser, 587–599, 917–929
- scavengers, 88–102
- Scelorchilus*, 1155–1160
- S. rubecula*, 515–525
- Scheiman, Daniel M., see Moore, Jeffrey E., et al.
- Schiffornis turdina*, 985–995
- S. turdinus*, 985–995
- Scoter, 950–962; White-winged, 950–962
- Screech-Owl, Eastern, 46–57, 738–751
- Scrub-Jay, Western, 1081–1093
- Scytalopus*, 752–766
- S. magellanicus*, 515–525
- seabirds, 1–6, 148–155, 380–390, 463–472, 752–766,
1161–1171
- Sealy, Spencer G., see Peer, Brian D., and —
- seasonal, — distributions, 610–618; — movements,
1005–1013; — patterns, 110–117
- sedentary birds, 682–694
- Sedgwick, James A., Site fidelity, territory fidelity,
and natal philopatry in Willow Flycatchers
(*Empidonax traillii*), 1103–1121
- Sedinger, James S., et al., Early environment
and recruitment of Black Brant (*Branta
bernica nigricans*) into the breeding population,
68–73
- Sedinger, James S., see Herzog, Mark P., and —
- Seiurus aurocapillus*, 726–737, 1040–1055, 1155–1160
- Selenidera culik*, 88–102
- Seleucidis melanoleuca*, 1155–1160
- self-medication, 1262–1268
- Sephanoides galeritus*, 515–525
- Serin, European, 302–307, 637–651
- Serinus canaria*, 184–191
- S. serinus*, 302–307, 637–651
- Serpophaga*, 415–421
- S. griseiceps*, 415–421
- S. munda*, 415–421
- S. subcristata*, 415–421
- Setophaga ruticilla*, 15–22, 788–795, 877–893, 1040–1055
- sex ratio, operational, 391–403
- sex-role reversal, 391–403
- sexual, — dimorphism, 178–183, 660–681, 788–795;
— selection, 178–183, 391–403, 788–795

- Shaffer, Terry L., A unified approach to analyzing nest success, 526-540
- Shaffer, Terry L., see Peak, Rebecca G., et al.
- Shag, 103-109; Rock, 243-252
- Shapiro, Leo H., et al., Reciprocal introgression between Golden-winged Warblers (*Vermivora chrysoptera*) and Blue-winged Warblers (*V. pinus*) in eastern North America, 1019-1030
- Shawkey, Matthew D., and Hill, Geoffrey E., Feathers at a fine scale, 652-655
- Shearwater, 422-434, 847-864; Audubon's, 847-864; Galapagos, 847-864; Little, 847-864; Little-Audubon's complex, 847-864; Newell's, 847-864; Tropical, 847-864
- Sheppardia, 23-34
- S. aequatorialis*, 23-34
- S. aurantiithorax*, 23-34
- S. cyornithopsis*, 23-34
- S. gunningi*, 23-34
- S. lowei*, 23-34
- S. montana*, 23-34
- S. sharpei*, 23-34
- Sheridan, Jennifer A., et al., Weak association between measures of health and reproductive success in Green-rumped Parrotlets (*Forpus passerinus*) in Venezuela, 717-725
- Sherman, Paul W., see Koenig, Walter D., and — shorebirds, 170-177, 333-344, 1094-1102
- Shoveler, Northern, 497-508, 587-599, 695-706
- Shrike, Brown, 35-45; Fiscal, 1226-1237; Lesser Gray, 557-564; Loggerhead, 1081-1093; Red-backed, 361-371
- Sialia*, 118-133
- S. currucoides*, 118-133, 1081-1093
- S. mexicana*, 118-133
- S. sialis*, 118-133, 302-307, 435-445, 652-655
- Sieving, Kathryn E., et al., Heterospecific facilitation of forest-boundary crossing by mobbing understory birds in north-central Florida, 738-751
- Sillett, T. Scott, see Jones, Jason, et al.
- Sillett, T. Scott, see Morrison, Scott A., et al.
- Sinclair, Pamela H., et al., Birds of the Yukon Territory (rev.), 982-983
- singing activity, 110-117
- Siphonorhis americana*, 985-995
- S. americanus*, 985-995
- S. brewsteri*, 985-995
- site selection, 796-805
- Sitta carolinensis*, 435-445
- S. europaea*, 1155-1160
- Sittasomus griseicapillus*, 88-102, 1155-1160
- Sittidae, 1155-1160
- size dimorphism, 391-403
- Skua, 243-252; Great, 170-177
- slough levees, 210-223
- Smith, Kimberly G., see Jahn, Alex E., et al.
- Smith, Ruth B., et al., Migratory movements of Sharpshinned Hawks (*Accipiter striatus*) captured in New Mexico in relation to prevalence, intensity, and biogeography of avian hematozoa, 837-846
- Smithornis capensis*, 1155-1160
- snow finches, 637-651
- social, — context, 110-117; — rank, 1070-1080
- Sockman, Keith W., see Hahn, Thomas P., et al.
- Somateria mollissima*, 333-344, 777-787
- song, 224-229; — characters, 224-229; — mimicry, 156-161; — production, 824-836
- songbirds, 726-737, 788-795, 824-836, 904-916, 1094-1102, 1155-1160, 1262-1268
- Sora, 1081-1093
- Sorenseon, Michael D., see Payne, Robert B., and — South Pacific birds, 637-651
- Southern Hemisphere, 1005-1013
- Spalding, Marilyn G., see Forrester, Donald J., and — Sparrow, 224-229, 1081-1093; American Tree, 276-277, 1238-1249; Ashy Rufous-crowned, 904-916; Bachman's, 74-87; Baird's, 1130-1145; Black-throated, 1081-1093; Brewer's, 1081-1093, 1269-1281; Chipping, 738-751; Clay-colored, 1130-1145; Field, 865-876; Fox, 162-169, 1269-1281; Grasshopper, 1130-1145; House, 118-133, 302-307, 837-846, 1187-1198, 1226-1237; Lark, 1081-1093; Lincoln's, 162-169; Mountain White-crowned, 1269-1281; Rufous-crowned, 446-451, 904-916; Savannah, 276-277, 473-485, 1081-1093, 1130-1145; Song, 391-403, 404-414, 652-655, 656-659, 738-751, 824-836, 877-893; Southern California Rufous-crowned, 904-916; Vesper, 1081-1093; White-crowned, 162-169, 276-277; White-throated, 162-169, 738-751
- Sparrowhawk, Black, 1226-1237; Eurasian, 1103-1121
- spatial, — analysis, 192-202; — interactions, 243-252; — redistribution, 380-390; — relationships, 192-202
- speciation, 682-694, 1014-1018
- species, — abundance, 515-525; — composition, 88-102; — distribution, 604-609; — identification, 824-836; — limits, 752-766; — richness, 88-102, 515-525, 604-609
- sperm competition, 473-485
- Spheniscidae, 148-155
- Sphenisciformes, 422-434
- Spheniscus*, 148-155, 422-434
- S. demersus*, 422-434
- S. humboldti*, 422-434
- S. magellanicus*, 148-155, 243-252, 422-434
- S. mendiculus*, 422-434
- Sphyrapius varius*, 509-514, 548-556
- Spiza*, 435-445
- S. americana*, 435-445, 767-776, 1250-1261
- Spizaetus melanoleucus*, 88-102
- S. ornatus*, 88-102
- Spizella*, 610-618
- S. altroglaris*, 610-618
- S. arborea*, 1238-1249
- S. breweri*, 610-618, 1081-1093, 1269-1281

- S. pallida*, 610–618, 1130–1145
S. passerina, 610–618, 738–751, 1081–1093
S. pusilla, 865–876
Spizixos semitorques, 435–445
Sporophila, 1005–1013
 spring, — arrival, 1269–1281; — condition hypothesis, 917–929
 stabilizing selection, 847–864
 stable-hydrogen isotope ratios, 837–846
 stable-isotope, — analyses, 333–344, 170–177;
 — ratios, 333–344
Stachyris poliocephala, 1155–1160
 standard avian calibration, 1–6
 Stanley, Thomas R., Estimating stage-specific daily survival probabilities of nests when nest age is unknown, 134–147
 Starling, 1122–1129; European, 110–117, 118–133, 372–379, 1187–1198
Stercorarius parasiticus, 652–655
S. skua, 170–177
Sterna eurygnatha, 243–252
S. hirundinacea, 243–252
S. maxima, 243–252
S. nilotica, 637–651
 Stettenheim, Peter, 265–266
 Stewart, Ian R. K., see Byers, Bruce E., et al.
 Stiles, F. Gary, and Altshuler, Douglas L., Conflicting terminology for wing measurements in ornithology and aerodynamics, 973–976
 Stokes, David L., see Boersma, P. Dee, et al.
 Stokes, David, review by, 1293–1296
 stomach contents, 148–155
 Stonechat, 796–805
 stopover, — ecology, 1040–1055; — habitats, 1040–1055
 Stotz, Douglas F., see Banks, Richard C., et al.
 Stover, Dollie M., see Shapiro, Leo H., et al.
Streptopelia risoria, 837–846
Streptoprocne zonaris, 88–102
 stress response, 276–277
 Strigidae, 35–45, 88–102
Strix, 35–45
S. aluco, 541–547, 1056–1069
S. occidentalis caurina, 806–810, 1056–1069
S. o. lucida, 1056–1069
S. o. occidentalis, 1056–1069
S. rufipes, 515–525
S. varia, 806–810
 Strong, Allan M., et al., Effect of prey biomass on reproductive success and mating strategy of Bicknell's Thrush (*Catharus bicknelli*), a polygynandrous songbird, 446–451
Struthidea cinerea, 1155–1160
Sturnella, 1081–1093
S. magna, 435–445, 767–776, 796–805, 1146–1154
S. neglecta, 134–147, 224–229, 824–836, 1130–1145
 Sturnidae, 1155–1160
Sturnus vulgaris, 110–117, 118–133, 372–379, 435–445, 1155–1160, 1187–1198
 Stutchbury, Bridget J. M., see Fedy, Bradley C., and — suboscines, 752–766, 1155–1160
 sunbird, double-collared, 23–34, 660–681
 Sunbird, Eastern Double-collared, 660–681; Greater Double-collared, 660–681; Loveridge's, 660–681; Moreau's, 660–681; Northern Double-collared, 660–681; Olive-bellied, 660–681; Regal, 660–681; Rufous-winged, 660–681; Southern Double-collared, 660–681; Tiny, 660–681
 superabundant food, 557–564
 supplemental feeding, 463–472
 Surfbird, 752–766
 survival, 68–73, 404–414; adult, 904–916, 1199–1207; adult male, 15–22; annual, 15–22, 904–916, 1199–1207; — estimates, 118–133; first year annual, 68–73; immediate, 1238–1249; juvenile, 894–903, 1146–1154, 1199–1207; nestling, 1214–1225; postfledgling, 452–462; — rates, 1056–1069; stage-specific daily, 134–147
 survivorship, postfledgling, 404–414
 Swallow, 88–102, 1122–1129; Bank, 985–995, 1187–1198; Barn, 777–787, 1122–1129; Cave, 1122–1129; Cliff, 1122–1129; Tree, 302–307, 345–353, 777–787, 1187–1198
 Sweeney, Robert J., et al., Evolutionary variation in feather waxes of passerine birds, 435–445
 Swift, 88–102; Common, 1187–1198; Pacific, 985–995; White-rumped, 985–995
 Swihart, Robert K., see Moore, Jeffrey E., et al.
Sylvia, 985–995
S. althaea, 985–995
S. atricapilla, 1155–1160
S. curruca, 985–995
S. marginalia, 985–995
S. minula, 985–995
 Sylviidae, 1155–1160
Sylviothorhynchus desmursii, 515–525
Synallaxis spixii, 752–766
 synchronization of fledging, 452–462
 synchronous nestling stages, 1226–1237
Syndactyla rufosuperciliata, 1155–1160
 synonymization, 415–421
Synthliboramphus, 985–995
S. hypoleucus, 1161–1171
Syristes sibilator, 88–102
 systematics, 637–651, 660–681, 752–766, 847–864
Tachycineta bicolor, 302–307, 345–353, 777–787, 1187–1198
Tachyphonus cristatus, 88–102
T. surinamus, 88–102
Tadorna ferruginea, 985–995
Taeniopygia guttata, 473–485
 Tapanelli, Robert, see Macías-Duarte, Alberto, et al.
 tag retention time, 1161–1171
 Takahashi, Akinori, see Deguchi, Tomohiro, et al.
 Tanager, 88–102; Scarlet, 1040–1055
Tangara cayana, 435–445

- T. chilensis*, 88–102
T. gyrola, 88–102
T. nigroviridis, 1155–1160
T. punctata, 88–102
T. varia, 88–102
T. vassorii, 1155–1160
T. velia, 88–102
 Tapaculos, 752–766; Chucao, 515–525; Magellanic, 515–525; Ochre-flanked, 515–525
 Tarof, Scott A., and Ratcliffe, Laurene M., Habitat characteristics and nest predation do not explain clustered breeding in Least Flycatchers (*Empidonax minimus*), 877–893
 Tavares, Erika S., et al., Phylogenetic relationships among some Neotropical parrot genera (Psittacidae) based on mitochondrial sequences, 230–242
 taxonomic, — changes, 415–421; — characters, traditional, 847–864; — validity, 415–421
 Teal, Blue-winged, 497–508, 526–540, 587–599, 695–706; Green-winged, 587–599
 temperate-zone breeders, 1238–1249
 temporal schedule, 203–209
 tension zone model, 565–586
Terenura, 88–102
T. spodiopila, 88–102
 terminology, 973–976
 Tern, 35–45; Cayenne, 243–252; Gull-billed, 637–651; Royal, 243–252; South American, 243–252
 territorial spatial structure, 877–893
 territory, — size, 1226–1237; — switching, 486–496
 testicular, — maturation, 473–485; — morphology, 473–485; — volume, 473–485
 Tetraoninae, 333–344
Thalassarche, 1–6
T. chrysostoma, 1208–1213
Thalurania furcata, 88–102
 Thamnophilidae, 1031–1039, 1155–1160
Thamnophilus, 1031–1039
T. amazonicus, 1031–1039
T. caerulescens, 1155–1160
T. divisorius, 1031–1039
T. doliatus, 1031–1039
T. insignis, 1031–1039
T. punctatus, 1031–1039
T. ruficapillus, 1155–1160
 thermodynamic costs, 777–787
 Thompson, Christopher W., and Kitaysky, Alexander S., Polymorphic flight-feather molt sequence in Tufted Puffins (*Fratercula cirrhata*): A rare phenomenon in birds, 35–45
 Thompson, Frank R., III, see Peak, Rebecca G., et al.
 Thornbill, Brown, 1199–1207
 Thrasher, Brown, 354–360, 738–751, 1103–1121
 Thraupidae, 88–102, 230–242
Thraupis episcopus, 88–102, 435–445
T. palmarum, 88–102
 Thrush, 88–102, 118–133, 1040–1055; Bicknell's, 446–451; Blue Rock, 985–995; Cocoa, 404–414; Creamy-bellied, 1005–1013; Gray-cheeked, 1040–1055; Hermit, 46–57, 162–169, 738–751, 1094–1102; Song, 1187–1198; Swainson's, 162–169, 1040–1055; Wood, 46–57, 354–360, 404–414, 726–737, 824–836, 865–876, 1040–1055
Thryomanes bewickii, 435–445
Thryothorus leucotis, 486–496
T. ludovicianus, 74–87, 738–751
Tiaris canora, 985–995
T. canorus, 985–995
T. olivacea, 985–995
T. olivaceus, 985–995
 time of day, 110–117
 timing, — of breeding, 345–353, 452–462; — of fledging, 463–472; — of nesting, 695–705
 Tit, 289–301; Bearded, 302–307; Blue, 345–353, 652–655; Great, 184–191, 354–360; Marsh, 404–414
 Titmouse, Tufted, 74–87, 738–751, 865–876
Tityra cayana, 88–102
Tockus damarensis, 565–586
T. erythrorhynchus damarensis, 565–586
T. erythrorhynchus rufirostris, 565–586
T. monteiri, 565–586
T. rufirostris, 565–586
Todirostrum pictum, 88–102
Tolmomyias assimilis, 88–102
T. poliocephalus, 88–102
 tongue movements, 509–514
 Tordoff, Harrison B., In Memoriam: Walter John Breckenridge, 1903–2003, 1286–1288
 toucans, 88–102
Touit costaricensis, 985–995
T. dilectissimus, 985–995
T. dilectissimus, 985–995
T. huetii, 88–102
T. purpurata, 88–102
 Tourenq, Christophe, et al., Spatial relationships between tree-nesting heron colonies and rice fields in the Camargue, France, 192–202
 Towhee, Collared, 1014–1018; Eastern, 74–87, 738–751, 824–836; Green-tailed, 289–301, 404–414, 1081–1093, 1269–1281; Spotted, 1014–1018, 1081–1093
Toxostoma curvirostre, 1081–1093
T. rufum, 354–360, 435–445, 738–751, 1103–1121
 Traylor, Joshua J., et al., Nesting ecology of White-winged Scoters (*Melanitta fusca deglandi*) at Redberry Lake, Saskatchewan, 950–962
Tringa totanus, 372–379, 877–893
 Trochilidae, 88–102
Troglodytes aedon, 302–307, 391–403, 738–751, 865–876, 1103–1121
T. troglodytes, 224–229, 435–445, 1155–1160
 Troglodytidae, 435–445
 Trogon, 88–102
T. melanurus, 88–102
T. violaceus, 88–102
T. viridis, 88–102

- Trogonidae, 88–102
 trogons, 88–102
 trophic organization, 88–102
 tropical dry forest, 1214–1225
 Turdidae, 23–34, 1155–1160
 turdids, 435–445
 Turdinae, 23–34, 88–102
Turdoides squamiceps, 1155–1160
Turdus albicollis, 88–102
T. amaurochalinus, 1005–1013
T. assimilis, 404–414
T. fumigatus, 404–414
T. merula, 1155–1160
T. migratorius, 162–169, 354–360, 404–414, 435–445, 738–751, 865–876, 1103–1121
T. olivaceus, 23–34
T. philomelos, 1155–1160, 1187–1198
T. solitarius, 985–995
 turkey, 717–725; domestic, 148–155
Turnagra capensis, 1155–1160
 Turnstone, Ruddy, 333–344
 Turtle-Dove, Ringed, 837–846
 Twinspot, Peters', 156–161
 type specimens, 415–421
Tyrannetes virescens, 88–102
 tyrannid, 752–766
 Tyrannidae, 88–102, 415–421, 1005–1013, 1155–1160
 Tyrannulet, 415–421; Gray-crowned, 415–421; White-bellied, 415–421
Tyrannulus elatus, 88–102
Tyrannus, 1081–1093
T. albogularis, 1005–1013
T. melancholicus, 88–102, 1005–1013, 1155–1160
T. savana, 88–102
T. tyrannus, 1103–1121
T. verticalis, 1081–1093
T. vociferans, 1081–1093
 Tyrant, 88–102; Patagonian, 515–525
Tyto alba, 103–109
 Tytonidae, 35–45

 undernourished nestlings, 1187–1198
 understory birds, 515–525, 738–751
 upland, 497–508
Upucerthia, 752–766
U. aricomae, 752–766
U. dumetaria, 752–766
U. validirostris, 752–766
Uraeginthus cyanocephalus, 156–161
 Urban, Emil K., In Memoriam: G. Stuart Keith, 1931–2003, 253–256
Uria, 148–155, 452–462
U. aalge, 35–45, 148–155, 1161–1171
U. lomvia, 148–155, 452–462
 uropygial wax, 435–445

 Valera, Francisco, see Hoi, Herbert, et al.
 Vangidae, 1155–1160

 Veery, 1040–1055
 vegetation, 1081–1093; — cover, 243–252;
 — structure, 894–903, 1130–1145
 vegetative, — characteristics, 1250–1261; — cover, 738–751
 Veit, M. Lisa, see Jones, Jason, et al.
Veniliornis cassini, 88–102
Vermivora, 610–618, 1014–1018
V. celata, 610–618, 1019–1030
V. chrysoptera, 1014–1018, 1019–1030, 1040–1055
V. peregrina, 435–445, 610–618, 1019–1030, 1040–1055
V. pinus, 1014–1018, 1019–1030, 1040–1055
V. ruficapilla, 610–618, 1019–1030, 1040–1055
 viability-predation hypothesis, 587–599
 vicariance, 682–694
 vicariant speciation, 682–694
 Vickery, Peter D., and Krohn, William B., In Memoriam: Ralph Simon Palmer, 1914–2003, 966–968
Vidua, 156–161
V. chalybeata, 156–161, 435–445
V. codringtoni, 156–161
V. obtusa, 156–161
V. paradisaea, 156–161
V. purpurascens, 156–161
 Villard, Pascal, and Cuisin, Jacques, How do woodpeckers extract grubs with their tongues? A study of the Guadeloupe Woodpecker (*Melanerpes herminieri*) in the French West Indies, 509–514
Vireo, 610–618
V. altiloquus, 88–102
V. bellii, 610–618
V. cassinii, 610–618
V. flavifrons, 738–751, 1040–1055
V. gilvus, 610–618, 1040–1055
V. griseus, 74–87, 610–618, 738–751
V. huttoni, 610–618
V. olivaceus, 88–102, 435–445, 738–751, 865–876, 1040–1055, 1094–1102, 1155–1160
V. o. chivi, 88–102
V. o. olivaceus, 88–102
V. philadelphicus, 1040–1055
Vireo, 1040–1055; Bell's, 610–618; Blue-headed, 1040–1055; Philadelphia, 1040–1055; Red-eyed, 738–751, 865–876, 1040–1055, 1094–1102; Warbling, 1040–1055; White-eyed, 74–87, 738–751; Yellow-throated, 738–751, 1040–1055
Vireolanus leucotis, 88–102
 Vireonidae, 88–102, 1155–1160
 vocal, — behavior, 184–191; — evolution, 224–229
 vocalizations, 184–191, 224–229, 486–496, 541–547, 565–586, 824–836
 Vogel, Uwe, see Göth, Ann, and —
 voice, 1031–1039
 Vuilleumier, François, In Memoriam: Jean Dorst, 1924–2001, 1289–1290
 vultures, 88–102, 837–846

- Wagner, Tansy L., see O'Connor, Raymond J., and —
 Wagtail, 682–694; Citrine, 777–787; Eastern Yellow, 985–995; Yellow, 985–995
- Walk, Jeffery W., et al., Renesting decisions and annual fecundity of female Dickcissels (*Spiza americana*) in Illinois, 1250–1261
- Walk, Jeffery W., see Kershner, Eric L., et al.
- Waller, Deborah A., see Revis, Hannah C., and —
- Warbler, 224–229, 877–893, 1014–1018, 1040–1055, 1238–1249; Bay-breasted, 162–169, 877–893, 1040–1055; Black-and-white, 738–751, 1040–1055; Blackburnian, 1040–1055; Blackpoll, 162–169, 1040–1055; Black-throated Blue, 15–22, 302–307, 473–485, 788–795, 1040–1055, 1094–1102; Black-throated Green, 162–169, 1040–1055; Blue-winged, 1014–1018, 1019–1030, 1040–1055; Brewster's, 1014–1018, 1019–1030; Bush, 1238–1249; Canada, 1040–1055; Cape May, 1040–1055; Cerulean, 7–14, 15–22; Chestnut-sided, 788–795, 1040–1055; Connecticut, 1040–1055; Golden-winged, 1014–1018, 1019–1030, 1040–1055; Great Reed, 203–209, 1226–1237; Hermit, 1014–1018; Hooded, 302–307, 404–414, 788–795, 796–805, 1040–1055, 1250–1261; Kentucky, 726–737, 1040–1055; Kirtland's, 877–893; Lawrence's, 1014–1018, 1019–1030; Magnolia, 162–169, 610–618, 1040–1055; Marsh, 203–209; Mourning, 1040–1055; Nashville, 1019–1030, 1040–1055; Orange-crowned, 1019–1030; Palm, 738–751; Pine, 74–87, 738–751; Prairie, 738–751, 796–805, 1040–1055, 1250–1261; Prothonotary, 354–360, 1103–1121; Reed, 203–209; Savi's, 35–45; Sedge, 203–209; Swainson's, 788–795; Tennessee, 1019–1030, 1040–1055; Townsend's, 15–22, 1014–1018; Willow, 302–307, 985–995, 1238–1249; Wilson's, 1040–1055; Worm-eating, 1040–1055; Yellow, 788–795, 1238–1249; Yellow-rumped, 15–22, 46–57, 74–87, 162–169, 289–301, 738–751, 1040–1055, 1094–1102; Yellow-throated, 1040–1055
- Ward, David H., see Sedinger, James S., et al.
- Warner, Richard E., see Kershner, Eric L., et al.
- Warner, Richard E., see Walk, Jeffery W., et al.
- Watanuki, Yutaka, see Deguchi, Tomohiro, et al.
- water levels, 894–903
- waterbirds, 192–202
- waterfowl, 35–45, 587–599, 695–706, 930–949, 1122–1129
- weather, 1056–1069, 1199–1207, 1269–1281
- Weaver, 1122–1129; Ploceine, 777–787; Sociable, 1199–1207; Village, 1172–1186
- Weaver, Heath B., and Brown, Charles R., Brood parasitism and egg transfer in Cave Swallows (*Parotchelidon fulva*) and Cliff Swallows (*P. pyrrhonota*) in south Texas, 1122–1129
- weaverbirds, 1172–1186, 1226–1237
- Wentworth, Kevin, see Walk, Jeffery W., et al.
- Westneat, David F., see Byers, Bruce E., et al.
- wetlands, 361–371, 497–508, 695–705
- white-blood-cell counts, 717–725
- White-eye, Japanese, 548–556
- Whitestart, 985–995
- Whitethroat, Desert, 985–995; Hume's, 985–995; Lesser, 985–995; Margelanica, 985–995
- Whitney, Bret M., et al., A new species of *Thamnophilus* antshrike (Aves: Thamnophilidae) from the Serra do Divisor, Acre, Brazil, 1031–1039
- Whydah, Broad-tailed Paradise, 156–161; Long-tailed Paradise, 156–161
- Widowbird, Jackson's, 1226–1237; Red-collared, 1226–1237
- Wigeon, American, 587–599
- Willson, Mary F., see Reid, Sharon, et al.
- Wilson, C. Morgan, and Holberton, Rebecca L., Individual risk versus immediate reproductive success: A basis for latitudinal differences in the adrenocortical response to stress in Yellow Warblers (*Dendroica petechia*), 1238–1249
- Wilsonia canadensis*, 1040–1055
- W. citrina*, 302–307, 404–414, 788–795, 796–805, 1040–1055, 1250–1261
- W. pusilla*, 1040–1055
- Wiltschko, Wolfgang, In Memoriam: Friedrich Wilhelm "Fritz" Merkel, 1911–2002, 1291
- wing, — length, 452–462; — measurements, 973–976
- Winkel, Wolfgang, see Bairlein, Franz, and —
- Winkler, David W., see Bowlin, Melissa S., and —
- Wins-Purdy, Andreas, see Drever, Mark C., et al.
- winter, — bird abundance, 46–57; — snowpack, 1269–1281
- wintering, — birds, 74–87; — grounds, 46–57
- Wiretail, Des Murs's, 515–525
- Wolf, Blair O., see Smith, Ruth B., et al.
- Woodcreeper, 88–102; Lesser, 1155–1160
- Woodhoopoe, Green, 178–183; Red-billed, 178–183
- Woodpecker, 58–67, 88–102, 178–183, 548–556; Acorn, 1081–1093; American Three-toed, 58–67; Black-backed, 58–67, 509–514; Downy, 74–87, 509–514, 738–751; Eurasian Three-toed, 58–67; Great Spotted, 58–67, 509–514; Green, 509–514; Guadeloupe, 509–514; Hairy, 509–514; Ladder-backed, 1081–1093; Lesser Spotted, 509–514; Middle Spotted, 58–67, 509–514; Pileated, 509–514; Red-bellied, 74–87, 738–751, 865–876; Red-cockaded, 58–67, 74–87, 1103–1121; Red-vented, 985–995; White-fronted, 548–556; Yucatan, 985–995
- Wood-Pewee, Eastern, 865–876, 1040–1055; Western, 824–836
- wood warblers, 110–117, 788–795
- Wrege, Peter H., see Emlen, Stephen T., and —
- Wren, 1155–1160, 1199–1207; Buff-breasted, 486–496; Cactus, 1081–1093; Carolina, 74–87, 738–751; House, 302–307, 391–403, 738–751, 865–876, 1103–1121; Marsh, 391–403, 1226–1237; South Island, 1155–1160; Winter, 224–229

- Xanthocephalus xanthocephalus*, 1226–1237
Xenicus gilviventris, 1155–1160
Xenops milleri, 88–102
Xipholena punicea, 88–102, 1155–1160
Xiphorhynchus pardalotus, 88–102
Xolmis coronata, 1005–1013
X. irrupero, 1155–1160
- Yamashita, Carlos, see Tavares, Erika S., et al.
Yellowthroat, Common, 738–751, 788–795, 865–876, 1040–1055
Yellow-Wagtail, Alaska, 985–995; Eastern, 985–995
yolk deposition, rapid, 103–109
Yorio, Pablo, see García Borboroglu, Pablo, and —
- Zenaida asiatica*, 1081–1093
Z. macoura, 354–360, 1081–1093, 1172–1186
Zhuravlev, Yury N., see Kulikova, Irina V., et al.
- Zimmerius gracilipes*, 88–102
Zimmerman, Guthrie S., see LaHaye, William S., et al.
Zink, Robert M., and Jones, Andrew W., review by, 1296–1298
Zonotrichia, 224–229
Z. albicollis, 162–169, 738–751
Z. atricapilla, 435–445
Z. leucophrys, 162–169
Z. l. oriantha, 1269–1281
Zoothera, 435–445
Z. citrina, 1155–1160
Z. interpres, 435–445
Z. piaggiae, 435–445
Zosteropidae, 1155–1160
Zosterops erythropleurus, 1155–1160
Z. japonicus, 548–556
Z. palpebrosa, 1155–1160

